

参考論文リスト

4. 環境濃度およびヒトへの曝露

- 4-1) Barton H. Predicted intake of trace elements and minerals via household drinking water by 6-year-old children from Krakow, Poland. Part 2: Cadmium, 1997-2001. *Food Addit Contam.* 2005 Sep;22(9):816-28.
- 4-2) Galal-Gorchev H. Dietary intake, levels in food and estimated intake of lead, cadmium, and mercury. *Food Addit Contam.* 1993 Jan-Feb;10(1):115-28. Review.
- 4-3) Lee JS, Chon HT, Kim KW. Human risk assessment of As, Cd, Cu and Zn in the abandoned metal mine site. *Environ Geochem Health.* 2005 Apr;27(2):185-91.
- 4-4) Louekari K. Estimation of dietary intake of cadmium: reliability of methods. *IARC Sci Publ.* 1992;(118):163-7. Review.
- 4-5) Peplow D, Edmonds R. Health risks associated with contamination of groundwater by abandoned mines near Twisp in Okanogan County, Washington, USA. *Environ Geochem Health.* 2004 Mar;26(1):69-79.
- 4-6) Puklova V, Batariova A, Cerna M, Kotlik B, Kratzer K, Melichercik J, Ruprich J, Rehurkova I, Spevakova V. Cadmium exposure pathways in the Czech urban population. *Cent Eur J Public Health.* 2005 Mar;13(1):11-9.

6. カドミウムによるヒトにおける有害性評価

- 6-1) WHO. Cadmium -Environmental Health Criteria 134-. Geneva, 1992.
- 6-2) Foulkes. Cadmium -Handbook of Experimental Pharmacology-. Berlin: Springer-Verlag, 1986.
- 6-3) 能勢隆之, 杉山恭子, 他. 銀ろう作業による急性カドミウム中毒の一例. *産業医学* 1988 ; 30 : 210-211.
- 6-4) 産業衛生学会. 許容濃度等の勧告(2002). *産衛誌*2002; 44:142.
- 6-5) Adams RG, Harrison JF, Scott P. The development of cadmium-induced proteinuria, impaired renal function and osteomalacia in alkaline battery workers. *Q J Med.* 1969; 38: 425-443.
- 6-6) Aoshima K. Environmental cadmium pollution and its health effects on inhabitants in Japan. Jinzu River basin: Clinical findings in Itai-itai disease. In, *Advances in the Prevention of Environmental Cadmium Pollution and Countermeasures*, 13-19, Nogawa K, Kurachi M, Kasuya M. (Eds.), Eiko Laboratory, Kanazawa, 1999.
- 6-7) 青島恵子 イタイイタイ病をめぐる最近の知見 *衛生化学*, 43:317-330、1997
- 6-8) Cai Y, Aoshima K, Katoh T, Teranishi H, Kasuya M. Renal tubular dysfunction in male inhabitants of a cadmium-polluted area in Toyama, Japan—an eleven-year follow-up study. *J Epidemiol.* 2001; 11: 180-189.
- 6-9) Ezaki T, Tsukahara T, Moriguchi j, Furuki K, Fukui Y, Ukai H, Okamoto S, Sakurai H, Honda S, Ikeda M. No clear-cut evidence for cadmium-induced renal tubular dysfunction among over

- 10,000 women in the Japanese general population: a nationwide large-scale survey *Int Arch Occup Environ Health*. 2003; 76: 186-196.
- 6-10) 樊 建軍、青島恵子、加藤輝隆、寺西秀豊、加須屋 実、富山県神通川流域カドミウム環境汚染地域住民の尿細管障害に関する追跡研究 第1報 土壌汚染改良事業開始後のカドミウム曝露の変化と尿細管障害の予後、*日衛誌*53:545-557、2000
- 6-11) Friberg L. Health hazards in the manufacture of alkaline accumulators with special reference to chronic cadmium poisoning. *Acta Med Scand*. 1950; Suppl. 240: 1-124.
- 6-12) Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Dietary exposure to cadmium at close to the current provisional tolerable weekly intake dose not affect renal function among female Japanese farmers. *Environ Res*. 2004; 95: 20-31.
- 6-13) Ikeda M, Zhang Z-W, Moon C-S, Shimbo S, Watanabe T, Nakatsuka H, Matsuda-Inoguchi N, Higashikawa K. Possible effects of environmental cadmium exposure on kidney function in the Japanese general population. *Int Arch Occup Environ Health* 2000; 73: 15-25.
- 6-14) Iwata K, Saito H, Moriyama M, Nakano A. Renal tubular function after reduction of environmental cadmium exposure: a ten-year follow-up. *Arch Environ Health*. 1993; 48: 157-163.
- 6-15) 神通川流域住民健康調査検討会報告書、平成15年7月
- 6-16) 「カドミウムによる環境汚染地域住民健康調査」検討委員会、カドミウムによる環境汚染地域住民健康調査、環境保健レポート、56:69-345、1989
- 6-17) カドミウム汚染地域住民健康影響調査検討会報告書、平成14年3月
- 6-18) Kazantzis G. Renal tubular dysfunction and abnormalities of calcium metabolism in cadmium workers. *Environ Health Perspect*. 1979; 28: 155-159.
- 6-19) Kido T, Honda R, Tsuritani I, Yamaya H, Ishizaki M, Yamada Y, Nogawa K. Progress of renal dysfunction in inhabitants environmentally exposed to cadmium. *Arch Environ Health*. 1988; 43: 213-217.
- 6-20) Nakadaira H, Nishi S. Effects of low-dose cadmium exposure on biological examinations. *Sci Total Environ*. 2003; 308: 49-62.
- 6-21) 村田 勇、イタイイタイ病の研究、*日本医師会雑誌*、65:15-42、1971
- 6-22) 斎藤 寛、中野篤浩、カドミウム環境汚染と人間の健康-生態中毒学のアプローチ “エコトキシコロジー” (大井 玄、鈴木継美、井村伸正編)、pp. 119-128、1983、篠原出版
- 6-23) Suwazono Y, Kobayashi E, Okubo Y, Nogawa K, Kido T, Nakagawa H. Renal effects of cadmium exposure in cadmium nonpolluted areas in Japan. *Environ Res*. 2000; 84: 44-55.
- 6-24) 武内重五郎、中本 安、イタイイタイ病、現代内科学大系 1969年刊追補、366-394、1969、中山書店
- 6-25) Tsuchiya, K. ed.: *Cadmium Studies in Japan: A Review*. pp. 144-252, Kodansha, 1978, Tokyo
- 6-26) 土屋健三郎、岩尾総一郎、秋田・石川・長崎3県のCd摂取量調査の結果と評価環境保健レポート、44:86-115、1978
- 6-27) World Health Organization (WHO), *Environmental Health Criteria 134: Cadmium*. pp. 174-192, World Health Organization, 1992, Geneva

- 6-28) Alfven T, Jarup L, Elinder C-G. Cadmium and lead in blood in relation to low bone mineral density and tubular proteinuria. *Environ Health Perspect*, 2002; 110: 699-702.
- 6-29) 青島恵子、加藤輝隆、寺西秀豊、堀口兵剛、加須屋 実、カドミウム腎症におけるカルシウム・リン・ビタミンD代謝異常－富山県神通川流域カドミウム汚染地域に見いだされた近位尿細管障害34例の検討－ *日衛誌* 1993; 47: 1009-1020.
- 6-30) 青島恵子、樊 建軍、加藤輝隆、寺西秀豊、加須屋 実、萩野茂継、カドミウム腎症における尿細管機能と骨代謝：15年間の追跡研究、*環境保健レポート* 2002; 68: 220-224.
- 6-31) Aoshima K, Fan J, Cai Y, Katoh T, Teranishi H, Kasuya M. Assessment of bone metabolism in cadmium-induced renal tubular dysfunction by measurements of biochemical markers. *Toxicol Lett*. 2003; 136: 183-192.
- 6-32) 原田孝司、原 耕平、緒方時雄、西村敬一、重野 哲、西村 昇、七條利幸、鋤塚 眞、佐藤佐由利、長崎県対馬カドミウム環境汚染地域における経過観察者の主要検査所見の推移、*環境保健レポート* 1991; 58: 205-211.
- 6-33) Honda R, Tsuritani I, Noborisaka Y, Suzuki H, Ishizaki M, Yamada Y. Urinary cadmium excretion is correlated with calcaneal bone mass in Japanese women living in an urban area. *Environ Res*. 2003; 91: 63-70.
- 6-34) Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Environmental exposure to cadmium at a level insufficient to induce renal tubular dysfunction does not affect bone density among female Japanese farmers. *Environ Res*. (in press)
- 6-35) 城戸照彦、大道正義、能川浩二、本多隆文、釣谷伊希子、石崎昌夫、山田裕一、北川正信、石川県梯川流域Cd汚染地住民（健康管理対象者）の一症例－臨床及び病理検査成績－、*環境保健レポート* 1991; 58: 161-165.
- 6-36) Kido T, Nogawa K, Yamada Y, Honda R, Tsuritani I, Ishizaki M, Yamaya H. Osteopenia in inhabitants with renal dysfunction induced by exposure to environmental cadmium. *Int Arch Occup Environ Health*. 1989; 61: 271-276.
- 6-37) Kido T, Honda R, Tsuritani I, Ishizaki M, Yamada Y, Nakagawa H, Nogawa K, Dohi Y. Serum levels of bone Gla-protein in inhabitants exposed to environmental cadmium. *Arch Environ Health*. 1991; 46: 43-49.
- 6-38) 村田 勇、イタイイタイ病の研究、*日本医師会雑誌* 1971; 65:15-42.
- 6-39) 中川昭忠、富山県に発生した骨軟化症の研究（所謂いたいたい病）、*金沢医理学叢書* 1960;56:1-51.
- 6-40) 中川秀昭、西条旨子、森河裕子、田畑正司、千間正美、三浦克之、由田克士、奥村義治、河野俊一、城戸照彦、北川正信、骨病理検索を行った梯川流域カドミウム汚染地域要管理者の1例、*環境保健レポート* 1993; 60: 130-135.
- 6-41) Nordberg G, Jin T, Bernard A, Fierens S, Buchet J P, Ye T, Kong Q, Wang H. Low bone density and renal dysfunction following environmental cadmium exposure in China. *Ambio*. 2002; 31: 478-481.
- 6-42) Takebayashi S, Jimi S, Segawa M, Kiyoshi Y. Cadmium induces osteomalacia mediated by proximal tubular atrophy and disturbances of phosphate reabsorption. A study of 11 autopsies. *Pathol Res Pract*. 2000; 196: 653-663.
- 6-43) Wu X, Jin T, Wang Z, Ye T, Kong Q, Nordberg G. Urinary calcium as a biomarker of renal

- dysfunction in a general population exposed to cadmium. *J Occup Environ Med.*, 2001; 43: 898-904.
- 6-44) Friberg L. Health hazards in the manufacture of alkaline accumulators with special reference to chronic cadmium. *Acta med Scand*, 1950; 138: 1-124.
- 6-45) Smith J P, Smith J C, McCall A J. Chronic poisoning from cadmium fume. *J Pathol Bacteriol*, 1960 ; 80 : 287-296.
- 6-46) Sakurai H, Omae K, Toyama T, et al, . Cross-sectional study of pulmonary function in cadmium alloy workers. *J Work Environ Health*, 1982 ; 8 : 122-130.
- 6-47) Tarasenko N, Vorobjava R. Hygienic problems connected with the use of cadmium. *Vestn Akad Med Nauk SSR*, 1973 ; 28 : 37-43.
- 6-48) Horstowa H, Sikorski M, Tyborski H. Chronic cadmium poisoning in the clinical and radiological picture. *Med Prac occup Med*, 1966 ; 17 : 13-25.
- 6-49) Lauwery R, Buchet J, Roela H, et.al, . Epidemiological survey of workers exposed to cadmium. *Arch environ Health*, 1974a ; 28 : 145-148.
- 6-50) Kjellstoem T. Exposure and accumulation of cadmium in populations from Japan, the United States, and Sweden *Environ. Health Perspect*, 1979 ; 28 : 169-197.
- 6-51) Armstrong B, Kazantzis G. The mortality of cadmium workers. *Lancet* 1983 ; 1 : 1424-1427.
- 6-52) Kazantzis G, Lam T, Sullivan K. Mortality of cadmium exposed workers: A five year update. *Scand J Work Environ Health*, 1988 ; 14 : 220-223.
- 6-53) Mannino D M, Holguin F, Savage-Brown A, et al, . Urinary cadmium levels predict lower lung function in current and former smokers : data from the Third National Health and Nutrition examination Survey. *Thorax* 2004 ; 59 : 194-198.
- 6-54) Schroeder HA, Vinton WH Jr. Hypertension induced in rats by small doses of cadmium. *Am J Physiol*, 1962, 202:515-518.
- 6-55) Perry HM Jr, Erlanger MW. Metal induced hypertension following chronic feeding of low dose of cadmium mercury. *L lab Clin Med*, 1974, 83:541-547.
- 6-56) Kopp ST, Glonek T, Perry HM Jr, Erlanger MW, Perry EF. Cardiovascular actions of cadmium at environmental exposure levels. *Science*, 1982, 217:837-838.
- 6-57) Dalhamm T, Friberg L. The effect of cadmium on blood pressure and respiration and the use of dimercaprol(BAL) as antidote. *Acta Pharmacol Toxicol*, 1954, 10:199-203.
- 6-58) Kopp SJ, Hawley PL. Factors influencing cadmium toxicity in A-V conduction system of isolated perfused rat heart. *Toxicol Appl Pharmacol*. 1976, 37:531-544.
- 6-59) Nogawa K and Kawano S A survey of the blood pressure of women suspected of itai-itai disease. *Juzen Med J* 1969 3:357-363.
- 6-60) The Japanese Public Health Association Cadmium Research Committee Studies of health effects of cadmium. *Kankyo Hoken Report* 1989 56:69-345.
- 6-61) Nishijo M, Nakagawa H, Kido T Environmental cadmium exposure and hypertension and cardiovascular risk. *Metal ions in biology and medicine*, vol.6, Centino JA, Colley P, Vernet G, Finkelman RB, Gibb H, Etienne JC (Eds), p635-637, ohn Libbery Eurotext, Paris, 2000.
- 6-62) Shinoda A, Yuri T, Nakagawa S Clinical findings of itai-itai disease patients. *Kankyo Hoken Report* 1977 41:44-52.

- 6-63) Kagamimori S, Naruse Y, Fujita T, Watanabe M, Nishino H, Shinmura T. Factors associated with blood pressure in females with heavy exposure to cadmium. *Bull Environ Contam Toxic* 1985 ; 35:356-392.
- 6-64) Aoshima K and Kasuya M. Environmental exposure to cadmium and effects on human health part3. The results of blood examination and blood pressure in inhabitants of the cadmium polluted Jinzu River basin in Toyama prefecture. *Jpn J Hyg* 1988 ; 43:949-955.
- 6-65) Kagamimori S, Watanabe M, Nakagawa H, Okumura Y, Nakagawa S. Case control study of cardiovascular function in females with a history of heavy exposure to cadmium. *Bull Environ Contam Toxic* 1986 ; 36:484-490.
- 6-66) Beevers DG, Campbell BC, Goldberg A, Moore MR, Hawthorne VM. Blood-cadmium in hypertensives and normotensives. *Lancet* 1976 ; 2: 222-224.
- 6-67) Staessen J, Bulpitt CJ, Roels H, Bernard A, Fagard R, Joossens JV. Urinary cadmium and lead concentrations and their relation to blood pressure in a population with low exposure. *Br J Ind Med* 1984 ; 41:241-248.
- 6-68) Staessen J, Amery A, Bernard A, Bruaux P, Buchet JP, Bulpitt CJ, Claeys F, De Plaen P, Ducoffre G, Fagard R, Lauwerys RR, Lijnen P, Nick L, Saint Remy A, Roels H, Rondia D, Sartor F, Thijs L. Blood pressure, the prevalence of cardiovascular diseases, and exposure to cadmium: a population study. *Am J Epidemiol* 1991 ; 134: 257-267.
- 6-69) Whittemore AS, DiCiccio Y, Provenzano G. Urinary cadmium and blood pressure: results from the NHANES II survey. *Environ Health Perspect* 1991 ; 91: 133-140.
- 6-70) International Agency for Research on Cancer. Cadmium and cadmium compounds. Monographs on evaluation of carcinogenic risks to humans 1993 ; 58 : 119-237.
- 6-71) Potts C. Cadmium proteinuria –The health of battery workers exposed to cadmium oxide dust. *Ann occup Hyg* 1965 ; 8 : 55-61.
- 6-72) Sorahan T, Waterhouse J. Mortality study of nickel-cadmium battery workers by the method of regression models in life tables. *Br J Ind Med* 1983 ; 40 : 293-300.
- 6-73) Sorahan T. Mortality from lung cancer among a cohort of nickel cadmium battery workers:1946-84. *Br J Ind Med* 1987 ; 44 : 803-809.
- 6-74) Sorahan T, Esmen NA. Lung cancer mortality in UK nickel-cadmium battery workers, 1947-2000. *Occup Environ Med* 2004 Feb ; 61(2) : 108-16.
- 6-75) Jarup L, Bellander T, Hogstedt C, et al,. Mortality and cancer incidence in Swedish battery workers exposed cadmium and nickel. *Occup Environ Med* 1998 ; 55 : 755-759.
- 6-76) Sorahan T, Lister A, Gilthorpe MS, et al,, Mortality of copper cadmium alloy workers with special reference to lung cancer and non-malignant disease of the respiratory system,1946-92 . *Occup Environ Med* 1995 ; 52 : 804-812.
- 6-77) Lemen R, Wagoner J, Blejer H. Cancer mortality among cadmium production workers. *Ann NY Acad Sci* 1976 ; 271 : 273-279.
- 6-78) Thun M, Schnorr T, Smith A, et al,. Mortality among a cohort of U.S.cadmium production workers-an update. *J Natl Cancer Inst* 1985 ; 74 : 325-333.
- 6-79) Stayner L, Smith R, Thun M, et al,. A quantitative assessment of lung cancer risk and occupational cadmium exposure. *IARC Sci Publ* 1992 ; 118 : 447-455.

- 6-80) Lamm S, Parkinson M, Anderson M, et al,. Determinants of lung cancer risk among cadmium exposed workers. *Ann Epidemiol* 1992 ; 2 : 195-211.
- 6-81) Sorahan T,Lancashire R. Lung cancer mortality in a cohort of workers employed at a cadmium recovery plant in the United states :an analysis with detailed job histories. *Occup Environ Med* 1997 ; 54 : 194-201.
- 6-82) Kazantzis G,Armstrong B. A mortality study of cadmium workers in the United Kingdom. *Stand J Work Environ Health* 1982 ; 8(1) : 157-160.
- 6-83) Armstrong B, Kazantzis G. The mortality of cadmium workers. *Lancet* 1983 ;1:1424-1427.
- 6-84) Kazantzis G, Lam T, Sullivan K. Mortality of cadmium exposed workers:A five year update. *Scand J Work Environ Health* 1988 ; 14 : 220-223.
- 6-85) Armstrong B, Kazantzis G. Prostatic cancer and chronic respiratory and renal disease in British cadmium workers :a case control study. *Br J ind Med* 1985 ; 42 : 540-545.
- 6-86) Ades A,Kazantzis G. Lung cancer in a non-ferrous smelter:the role of cadmium. *Br J ind Med* 1988 ; 45 : 435-442.
- 6-87) Morgan J. Cadmium and zinc abnormalities in bronchogenic carcinoma. *Cancer* 1970 ; 25 : 1394-1398.
- 6-88) Morgan J, Burch H, Watkiss J. Tissue cadmium and zinc content in emphysema and bronchogenic carcinoma. *J chron Dis* 1971 ; 24 : 107-110.
- 6-89) Shigematsu I, Morita M, Nagai M,et al,. A retrospective mortality study on cadmium -exposed pollution in Japan: An investigation on mortality rates by a pollution level in Toyama Prefecture. *Kankyo hoken Rep* 1982b ; 48(suppl) : 118-136.
- 6-90) Arisawa K, Nakano A, Saito H, et al,. Mortality and cancer incidence among a population previously exposed to environmental cadmium. *Int Arch Occup Environ Health* 2001 ; 74 : 255-262.
- 6-91) Nishijo M, Nakagawa H, Tabata M. Environmental expose level and blood and urinary cadmium concentrarions. *Environ Sci* 1995 ; 3 : 125-135.
- 6-92) Nakagawa H, Nishijo M, Morikawa Y, et al,. Increased urinary beta2-microgroblin and mortality rate by cause of death in a cadmium- polluted area. *Environ Health Prevent Med* 1996 ; 1 : 144-148.
- 6-93) 重松逸造, 竹内重五郎, 簗輪眞澄, 永井正規, 喜田村正次, 臼井竹次郎, 福島匡昭. カドミウム汚染地域住民の死因に関する疫学調査研究. *環境保健レポート* 1980 46 (Part 2):1-71.
- 6-94) Kawano, S., Nakagawa, H., Okumura Y, Tsujikawa K. A mortality study of patientswith itai-itai disease. *Environ Res* 1986 40:98-102.
- 6-95) Nakagawa, H., Tabata M, Morikawa, Y., Senma M, Kitagawa Y, Kawano, S., Kido T. High mortality and shortened life-span in patients with itai-itai disease and subjects with suspected disease. *Arch Environ Health* 1991 45:283-287.
- 6-96) Matsuda T, Kobayashi E, Okubo Y, Suwazono Y, Kido T. Nishijo M, Nakagawa H, Nogawa K Assosiation between renal dysfunction and mortality among inhabitants in the region around Jinzu River basin polluted by cadmium. *Environ. Res.* 2002 88:156-163
- 6-97) Ishihara T, Kobayashi E, Okubo Y, Suwazono Y, Kido T. Nishijo M, Nakagawa H, Nogawa K

- Assosiation between cadmium concentration in rice and mortality in the Jinzu River basin, Japan. *Toxicology* 2001 163:23-28.
- 6-98) Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nishijo M, Nakagawa H, Nogawa K
Assosiation between total cadmium intake calculated from the cadmium concentration in household rice and mortality among inhabitants of the cadmium-polluted Jinzu River basin of Japan. *Toxicology Lett* 2002 129:85-91.
- 6-99) Nakagawa H., Kawano, S., Okumura T, Fujita T., Nishi, M Mortality study of inhabitants in a cadmium-polluted area. *Bull Environ Contam Toxicol* 1987 38:553-60.
- 6-100) Nishijo, M., Nakagawa, H Morikawa, Y., Tabata, M., Senma M., Miura, K., Takahara H., Kawano, S., Nishi, M., Mizukoshi, K., Kido, T., Nogawa, K., 1995. Mortality of inhabitants in an area polluted by cadmium: 15 year follow up. *Occup. Environ. Med.* 2, 181-184.
- 6-101) Nakagawa, H., Nishijo, M., Morikawa, Y., Tabata, M., Senma, M., Kitagawa, Y., Kawano, S., Sugita, N., Nishi, M., Kido, T., Nogawa, K., 1993. Urinary $\beta 2$ - microglobulin concentration and mortality in a cadmium-polluted area. *Arch. Environ. Health* 48, 428-435.
- 6-102) Nishijo, M., Nakagawa, H Morikawa, Y., Tabata, M., Senma, M., Kitagawa, Y., Kawano, S., Sugita, N., Nishi, M., Kido, T., Nogawa, K., 1994. Prognostic factors of renal dysfunction induced by environmental cadmium pollution. *Environ. Res.* 64, 112-121.
- 6-103) Nakagawa, H., Nishijo, M., Morikawa, Y., Tabata, M., Miura, K., Takahara, H., Okumura, Y., Yoshita, K., Kawano, K., Nishi, M., Kido, T., Nogawa K. (1996b) Increased urinary $\beta 2$ -microglobulin and mortality rate by cause of death in a cadmium-polluted area. *Environ. Health Prev. Med.* 1, 144-148.
- 6-104) 中川秀昭：カドミウム汚染地域住民の健康障害に関する研究 腎尿細管障害程度およびカドミウム暴露量と生命予後－15年間の追跡調査－、環境保健レポート、1999、65:76-79.
- 6-105) Iwata K, Saito H, Moriyama M, Nakano A Association between renal tubular dysfunction and mortality among residents in a cadmium polluted area, Nagasaki Japan. *Tohoku J Exp Med* 1991 164:93-102.
- 6-106) Iwata K, Saito H, Nakano A Association between renal tubular dysfunction and mortality: further evidence. *Tohoku J Exp Med* 1991 164:319-330.
- 6-107) Iwata, K., Saito, H., Moriyama, M., Nakano, A. Follow up study of renal tubular dysfunction and mortality in residents of an area polluted with cadmium. *Br J Ind Med* ,1992 49:736-737.
- 6-108) Arisawa K, Nakano A, Saito H, Liu XJ, Yokoo M, Soda M, Koba T, Takahashi T, Kinoshita K. Mortality and cancer incidence among a population previously exposed to environmental cadmium. *Int Arch Occup Environ Health* 2001 74:255-262.
- 6-109) 小山 洋, 佐藤雅彦, 遠山千春 低容量カドミウム暴露と健康影響(4) 生体負荷量・臨界濃度・生命予後。(2003) 日本衛生学雑誌、57, 624-635.
- 6-110) Urinary cadmium levels predict lower lung function in current and former smokers: data from the Third National Health and Nutrition Examination Survey. Mannino DM, Holguin F, Greves HM, Savage-Brown A, Stock AL, Jones RL. *Thorax*. 2004 Mar; 59 (3); 194 -198.
- 6-111) Lung cancer mortality in UK nickel-cadmium battery workers, 1947 – 2000. Sorahan T, Esmen

- NA. *Occup Environ Med*. 2004 Feb; 61 (2): 108 – 116.
- 6-112) Renal effects of cadmium exposure in cadmium nonpolluted areas in Japan. Suwazono Y, Kobayashi E, Okubo Y, Nogawa K, Kido T, Nakagawa H. *Environ Res*. 2000 Sep; 84(1): 44-55.
- 6-113) No clear-cut evidence for cadmium-induced renal tubular dysfunction among over 10,000 women in the Japanese general population: a nationwide large-scale survey. Ezaki T, Tsukahara T, Moriguchi J, Furuki K, Fukui Y, Ukai H, Okamoto S, Sakurai H, Honda S, Ikeda M. *Int. Arch. Occup. Environ. Health*. 2003 Apr; 76(3): 186-196.
- 6-114) Dietary exposure to cadmium at close to the current provisional tolerable weekly intake dose not affect renal function among female Japanese farmers. Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. *Environ Res*. 2004 May; 95 (1): 20-31.
- 6-115) Pharmacokinetics/pharmacodynamics (PK/PD) modeling of risks of kidney toxicity from exposure to cadmium: estimates of dietary risks in the U.S. population. Diamond GL, Thayer WC, Choudhury H. *J Toxicol Environ Health A*. 2003 Nov 28; 66(22): 2141-2164.
- 6-116) Adams RG. Clinical and biochemical observation in men with cadmium nephropathy. A twenty-year study. *Arh Hig Rada Toksikol*. 1979;30:219-31.
- 6-117) Adams RG, Harrison JF, Scott P. The development of cadmium-induced proteinuria, impaired renal function, and osteomalacia in alkaline battery workers. *Q J Med*. 1969 Oct;38(152):425-43.
- 6-118) Baader EW. Chronic cadmium poisoning. *Disch, Med Wochenschr*. 1951;76:484-7.
- 6-119) Bernard A, Buchet JP, Roels H, Masson P, Lauwerys R. Renal excretion of proteins and enzymes in workers exposed to cadmium. *Eur J Clin Invest*. 1979 Feb;9(1):11-22.
- 6-120) Bonnell JA. Emphysema and proteinuria in men casting copper-cadmium alloys. *Br J Ind Med*. 1955;12:181-97.
- 6-121) Bonnell JA, Kazantzis G, King EA. A follow-up study of men exposed to cadmium oxide fume. *Br J Ind Med*. 1960;17:69-80.
- 6-122) De Silva PE, Donnan MB. Chronic cadmium poisoning in a pigment manufacturing plant. *Br J Ind Med*. 1981 Feb;38(1):76-86.
- 6-123) Elinder CG, Edling C, Lindberg E, Kagedal B, Vesterberg O. beta 2-Microglobulinuria among workers previously exposed to cadmium: follow-up and dose-response analyses. *Am J Ind Med*. 1985;8(6):553-64.
- 6-124) Friberg L. Health hazards in the manufacture of alkaline accumulators with special reference to chronic cadmium poisoning. Doctorial thesis. *Acta Med Scand* 1950;138(s240):1-124.
- 6-125) Friberg L. Proteinuria and emphysema among workers exposed to cadmium and nickel dust in a storage battery plant. *Proc Int Congr Ind Med* 1948a; 9:641-4.
- 6-126) Friberg L. Proteinuria and kidney injury among workmen exposed to cadmium and nickel dust. *J Ind Hyg Toxicol* 1984b; 30:32-6.

- 6-127) Hansen L, Kjellström T, Vesterberg O. Evaluation of different urinary proteins excreted after Cd exposure. *Int Arch Occup Environ Health*. 1977;40:273-82
- 6-128) Kazantzis G, Flynn FV, Spowage JS, Trott DG. Renal tubular malfunction and pulmonary emphysema in cadmium pigment workers. *Q J Med*. 1963 Apr;32:165-192.
- 6-129) Kawada T, Shinmyo RR, Suzuki S. Changes in urinary cadmium excretion among pigment workers with improvement of the work environment. *Ind Health*. 1993;31(4):165-70.
- 6-130) Kjellstrom T, Friberg L, Rahnster B. Mortality and cancer morbidity among cadmium-exposed workers. *Environ Health Perspect*. 1979 Feb;28:199-204.
- 6-131) Lauwerys RR, Buchet JP, Roels HA, Brouwers J, Stanescu D. Epidemiological survey of workers exposed to cadmium. *Arch Environ Health*. 1974 Mar;28(3):145-8.
- 6-132) Lauwerys RR, Roels HA, Buchet JP, Bernard A, Stanescu D. Investigations on the lung and kidney function in workers exposed to cadmium. *Environ Health Perspect*. 1979 Feb;28:137-45.
- 6-133) Lemen RA, Lee JS, Wagoner JK, Blejer HP. Cancer mortality among cadmium production workers. *Ann N Y Acad Sci*. 1976;271:273-9.
- 6-134) McDiarmid MA, Freeman CS, Grossman EA, Martonik J. Follow-up of biologic monitoring results in cadmium workers removed from exposure. *Am J Ind Med*. 1997 Sep;32(3):261-7.
- 6-135) NicaudP, LafitteA, Gross A. Symptoms of chronic cadmium intoxication. *Arch Mal Prof Med Trv Secur Soc*. 1942;4:192-202.
- 6-136) Sakurai H, Omae K, Toyama T, Higashi T, Nakadate T. Cross-sectional study of pulmonary function in cadmium alloy workers. *Scand J Work Environ Health*. 1982;8 Suppl 1:122-30.
- 6-137) Scott R, Paterson PJ, Mills EA, McKirdy A, Fell GS, Ottoway JM, Husain FE, Fitzgerald -Finch OP, Yates AJ, Lamont A, Roxburgh S. Clinical and biochemical abnormalities in coppersmiths exposed to cadmium. *Lancet*. 1976 Aug 21;2(7982):396-8.
- 6-138) Suzuki Y, Suzuki T, Ashizawa M. Proteinuria due to inhalation of cadmium stearate dust. *Ind Health*. 1965;3:73-85.
- 6-139) Tuchiya K. Proteinuria of workers exposed to cadmium fume. The relationship to concentration in the working environment. *Arch Environ Health*. 1967;14:875-80.
- 6-140) Thun MJ, Schnorr TM, Smith AB, Halperin WE, Lemen RA. Mortality among a cohort of U.S. cadmium production workers--an update. *J Natl Cancer Inst*. 1985 Feb;74(2):325-33.
- 6-141) Vorob'eva RS, Eremeeva EP. [State of the cardiovascular system in cadmium exposure] *Gig Sanit*. 1980 Oct;(10):22-5.
- 6-142) Aoshima K. Epidemiology of renal tubular dysfunction in the inhabitants of a Cadmium -polluted area in the Jinzu River basin in Toyama prefecture. *Tohoku J Exp Med*. 152, 151-172, 1987.
- 6-143) 平成 16 年度版環境白書、環境省
- 6-144) 石崎有信、福島匡昭。イタイイタイ病。 *日衛誌* 23, 271-285, 1968.
- 6-145) 福島匡昭、石崎有信、坂元倫子、能川浩二、小林悦子。 イタイイタイ病発生住民の腎障害に関する疫学的研究（第 1 報）。神通川流域住民の尿検査成績。 *日本公衛誌* 21, 65-73, 1974.
- 6-146) 福島匡昭、石崎有信、坂元倫子、能川浩二、小林悦子。 イタイイタイ病発生住民の腎

- 障害に関する疫学的研究（第2報）。精検者の尿異常所見と Cd 排泄の居住地および診断との関係に関する観察。日本公衛誌 22, 217-224, 1975.
- 6-147) 福山裕三、城石和子、久保田憲太郎. イタイイタイ病検診における尿蛋白のスクリーニング法の検討。医学と生物学 83, 85-90, 1971a.
- 6-148) 福山裕三. カドミウム汚染にともなう尿蛋白と各種腎疾患の尿蛋白の鑑別の試み。医学と生物学 83, 139-143, 1971b.
- 6-149) 福山裕三、川辺外美枝、城石和子、久保田憲太郎. イタイイタイ病発生地域住民の尿中カドミウム濃度。医学と生物学 85, 257-261, 1972.
- 6-150) 金井正光、野本昭三、笹岡澄子、内貴正治. イタイイタイ病における尿中 RBP 排泄とその臨床的意義。臨床化学シンポジウム 11, 194-199, 1971.
- 6-151) 金井正光、百瀬正、相沢孝夫、奥平貞英、本間達二、萩野昇. イタイイタイ病患者とその家族における尿中および血中 β_2 -microglobulin 測定の臨床的意義について。最新医学 31, 1793-1799, 1976.
- 6-152) Kanai M, Iwanaga T, Hagino N, Muto Y. Retinol-binding protein in tubular proteinuria of patients with Itai-Itai disease. *Wld Rev Nutr Diet* 31, 31-36, 1978.
- 6-153) 小林悦子. 環境中カドミウムの人体影響に関する疫学的研究（第1報）。性、年齢別尿検査成績。日本公衛誌 29, 123-133, 1982a.
- 6-154) 小林悦子. 環境中カドミウムの人体影響に関する疫学的研究（第2報）。Cd 汚染地居住期間別尿所見。日本公衛誌 29, 201-207, 1982b.
- 6-155) 能川浩二、小林悦子、本多隆文、石崎有信、河野俊一、大村利志隆、中川秀昭、梶博久、松田晴夫. 慢性カドミウム中毒の臨床生化学的研究（第5報）腎機能。日衛誌 36, 512-517, 1981.
- 6-156) Kjellström T, Shiroishi K, Evrin PE. Urinary β_2 -microglobulin excretion among people exposed to cadmium in the general environment: an epidemiological study in cooperation between Japan and Sweden. *Environ Res* 13, 318-344, 1977.
- 6-157) Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nogawa. Dose-response relationship between total cadmium intake calculated from the cadmium concentration in rice collected from each household of farmers and renal dysfunction in inhabitants of the Jinzu River basin, Japan. *J Appl Toxicol* 22, 431-436, 2002.
- 6-158) Nogawa K, Kobayashi E, Inaoka H, Ishizaki A. The relationship between the renal effects of cadmium and cadmium concentration in urine among the inhabitants of cadmium-polluted areas. *Environ Res* 14, 391-400, 1977.
- 6-159) Nogawa K, Ishizaki A, Kobayashi E. A comparison between health effects of cadmium and cadmium concentration in urine among inhabitants of the Itai-iai disease endemic district. *Environ Res* 18, 397-409, 1979a.
- 6-160) Nogawa K, Kobayashi E, Honda R. A study of the relation ship between cadmium concentrations in urine and renal effects of cadmium. *Environ Health Perspect* 28, 161-168, 1979b.
- 6-161) Nogawa K, Kobayashi E, Honda R, Ishizaki A, Kawano S, Matsuda H. Renal dysfunction of inhabitants in a cadmium-polluted area. *Environ Res* 23, 13-23, 1980.
- 6-162) Nogawa K, Yamada Y, Honda R, Tsuritani I, Ishizaki M, Sakamoto M. Urinary N-acetyl- β

- D-glucosaminidase and β_2 -microglobulin in 'itai-itai' disease. *Toxicol Lett* 16, 317-322, 1983.
- 6-163) Nogawa K. Biological indicators of cadmium nephrotoxicity in persons with low-level cadmium exposure. *Environ Health Perspect* 54, 163-169, 1984.
- 6-164) Osawa T, Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nogawa K. Retrospective study on the relationship between renal dysfunction and cadmium concentration in rice in individual hamlets in the Jinzu River basin, Toyama prefecture, Japan. *Environ Res* 86, 51-59, 2001.
- 6-165) Chiyoda N, Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nogawa K. Allowable level of lifetime cadmium intake calculated from the individuals in the Jinzu River basin of Japan. *Biolog Trace Element Res* 96, 9-20, 2003.
- 6-166) 城石和子、穴山道子、谷井美樹子、福山裕三、久保田憲太郎. イタイイタイ病発生地域住民の年齢別尿所見. *医学と生物学* 85, 263-267, 1972.
- 6-167) Shiroishi K, Kjellström T, Kubota K, Evrin PE, Anayama M, Vesterberg O, Shimada T, Piscator M, Iwata T, Nishino H. Urine analysis for detection of cadmium-induced renal changes, with special reference to β_2 -microglobulin: a cooperative study between Japan and Sweden. *Environ Res* 13, 407-427, 1977.
- 6-168) Watanabe K, Kobayashi E, Suwazono Y, Okubo Y, Kido T, Nogawa K. Tolerable lifetime cadmium intake calculated from the inhabitants living in the Jinzu River basin, Japan. *Bull Environ Contam Toxicol* 72, 1091-1097, 2004.
- 6-169) Watanabe Y, Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nogawa K. Relationship between cadmium concentration in rice and renal dysfunction in individual subjects of the Jinzu River basin determined using a logistic regression analysis. *Toxicology* 172, 93-101, 2002.
- 6-170) Nogawa K, Yamada Y, Kido T, Honda R, Ishizaki M, Tsuritani I, Kobayashi E. Significance of elevated urinary *N*-acetyl- β -D-glucosaminidase activity in chronic cadmium poisoning. *Sci Total Environ* 53, 173-178, 1986.
- 6-171) Tohyama C, Shaikh Z, Nogawa K, Kobayashi E, Honda R. Elevated urinary excretion of metallothionein due to environmental cadmium exposure. *Toxicology* 20, 289-297, 1981.
- 6-172) Tohyama C, Shaikh Z, Nogawa K, Kobayashi E, Honda R. Urinary metallothionein as a new index of renal dysfunction in "itai-itai" disease patients and other Japanese women environmentally exposed to cadmium. *Arch Toxicol* 50, 159-166, 1982.
- 6-173) Hayano M, Nogawa K, Kido T, Kobayashi E, Honda R, Tsuritani I. Dose-response relationships between urinary cadmium concentration and β -microglobulinuria using logistic regression analysis. *Arch Environ Health*. 1996 March/April;51(2):162-7.
- 6-174) Hoshi Y, Kido T, Nogawa K, Kito H, Shaikh ZA. Dose-response relationship between cadmium intake and prevalence of renal dysfunction using general linear models. *J Appl Toxicology*. 1995 Feb; 15(2):109-16.
- 6-175) Ishizaki M, Kido T, Honda R, Tsuritani I, Yamada Y, Nakagawa H, Nogawa K. Dose-response relationship between urinary cadmium and beta 2-microglobulin in a Japanese environmentally cadmium exposed population. *Toxicology*. 1989 Oct 2;58(2):121-31.
- 6-176) Kido T, Honda R, Tsuritani I, Ishizaki M, Yamada Y, Nogawa K. Progress of renal dysfunction in inhabitants environmentally exposed to cadmium. *Arch Environ Health*. 1988 May/June;43(3):213-7.

- 6-177) Kido T, Honda R, Tsuritani I, Yamaya H, Ishizaki M, Yamada Y, Nogawa K. [An epidemiological study on renal dysfunction of inhabitants in Cd-exposed areas in the Kakehashi River basin in Ishikawa Prefecture] *Nippon Eiseigaku Zasshi*. 1987 Dec; 42 (5):964-72.
- 6-178) Kido T, Nogawa K, Ishizaki M, Honda R, Tsuritani I, Yamada Y, Nakagawa H, Nishi M. Long-term observation of serum creatinine and arterial blood pH in persons with cadmium-induced renal dysfunction. *Arch Environ Health*. 1990 Jan/Feb;45(1):35-41.
- 6-179) Kido T, Shaikh ZA, Kito H, Honda R, Nogawa K. Dose-response relationship between urinary cadmium and metallothionein in a Japanese population environmentally exposed to cadmium. *Toxicology*. 1991 Jan;65(3):325-32.
- 6-180) Nakashima K, Kobayashi E, Nogawa K, Kido T, Honda R. Concentration of cadmium in rice and urinary indicators of renal dysfunction. *Occup Environ Med*. 1997 Oct;54(10):750-5.
- 6-181) Nishijo M, Nakagawa H, Morikawa Y, Tabata M, Miura K, Yoshita K, Higashiguchi K, Seto T, Kido T, Nogawa K, Mizukoshi K, Nishi M. Relationship between urinary cadmium and mortality among inhabitants living in a cadmium polluted area in Japan. *Toxico Lett*. 1999 Sep;108(2-3):321-7.
- 6-182) Nogawa K, Honda R, Kido T, Tsuritani I, Yamada Y, Ishizaki M, Yamaya H. A dose-response analysis of cadmium in the general environment with special reference to total cadmium intake limit. *Environ Res*. 1989 Feb;48(1):7-16.
- 6-183) Nogawa K, Ishizaki A, Kawano S. Statistical observations of the dose-response relationships of cadmium based on epidemiological studies in the Kakehashi River basin. *Environ Res*. 1978 Apr;15(2):185-98.
- 6-184) 斎藤 寛, 塩路隆治, 古川洋太郎, 有川 卓, 斎藤喬雄, 永井謙一, 道又勇一, 佐々木康彦, 古山 隆, 吉永 馨. カドミウム環境汚染にもとづく慢性カドミウム中毒の研究 秋田県小坂町細越地域住民に多発したカドミウムによる腎障害(多発性近位尿細管機能異常症)について. *日内会誌* 64, 37-49, 1975.
- 6-185) Saito H, Shioji R, Hurukawa Y, Nagai K, Arikawa T, Saito T, Sasaki Y, Furuyama T, Yoshinaga K. Cadmium-induced proximal tubular dysfunction in a cadmium-polluted area. *Contr Nephrol* 6, 1-12, 1977a.
- 6-186) 斎藤 寛, 永井謙一, 有川 卓, 斎藤喬雄, 塩路隆治, 古川洋太郎, 古山 隆, 吉永 馨. カドミウム環境汚染地域住民の尿 β 2-microglobulin 濃度-カドミウム負荷量との Dose-Effect Relationship について. *医学のあゆみ*, 100, 350-352, 1977b.
- 6-187) 蓓 幸三, 斎藤 寛, 中野篤浩, 海上 寛, 高田健右, 佐藤徳太郎, 古山 隆, 吉永 馨, 有川 卓, 永井謙一. カドミウム環境汚染地域住民の尿中 β 2-microglobulin, 一世代別, 性別の検討, ならびに近位尿細管検査成績との比較. *日腎誌* 23, 45-62, 1981.
- 6-188) Kojima S, Haga Y, Kurihara T, Yamawaki T, Kjellstrom T. A comparison between cadmium and urinary β 2-Microglobulin, total protein, and cadmium among Japanese farmers. *Environ Res* 14, 436-451, 1977.
- 6-189) 小野雅司, 斎藤 寛. 秋田県小坂町住民の死亡原因に関する疫学的研究. *日衛誌* 40, 799-811, 1985.
- 6-190) Iwata K, Saito H, Moriyama M, Nakano A. Follow-up study of renal tubular dysfunction and

- mortality in residents of an area polluted with cadmium. *Br J Ind Med* 49, 736-737, 1992.
- 6-191) 中野篤浩, 斎藤 寛, 脇阪一郎. カドミウム土壤汚染地域住民におけるカドミウムとβ2-マイクログロブリンの尿中排泄に関する研究. 国立公害研究所研究報告, 84, 13-30, 1985.
- 6-192) 小林悦子, 杉平直子, 中野篤浩, 遠山千春, 三種裕子, 斎藤 寛, 脇阪一郎. 長崎県対馬カドミウム汚染地住民における血液検査成績. 国立公害研究所研究報告, 84, 37-45, 1985.
- 6-193) Tohyama C, Kobayashi E, Saito H, Sugihara N, Nakano A, Mitane Y. Urinary α1-microglobulin as an indicator protein of renal tubular dysfunction caused by environmental cadmium exposure. *J Appl Toxicol* 6, 171-178, 1986.
- 6-194) Tohyama C, Mitane Y, Kobayashi E, Sugihara N, Nakano A, Saito H. The relationships of urinary metallothionein with other indicators of renal dysfunction in people living in a cadmium-polluted area in Japan. *J Appl Toxicol* 8, 15-21, 1988.
- 6-195) Iwata K, Saito H, Moriyama M, Nakano A. Renal tubular function after reduction of environmental cadmium exposure: A ten-year follow-up. *Arch Environ Health* 48, 157-163, 1993.
- 6-196) 劉曉潔. 長崎県対馬カドミウム土壤汚染地域住民の頭髮, 尿および血液カドミウム濃度—土壤復元前後 18 年での比較—. *日衛誌*, 54, 544-551, 1999.
- 6-197) Liu X-J, Arisawa K, Nakano A, Saito H, Takahashi T, Kosaka A. Significance of cadmium concentrations in blood and hair as an indicator of dose 15 years after the reduction of environmental exposure to cadmium. *Toxicol Lett* 123, 135-141, 2001.
- 6-198) 原田孝司, 平井義修, 原耕平, 嘉村末男. カドミウム環境汚染地域における経過観察者の近位尿細管障害の推移. *環境保健レポート* 1988; 54, 127-33.
- 6-199) Iwata K, Saito H, Moriyama M, Nakano A. Association between renal tubular dysfunction and mortality among residents in a cadmium-polluted area, Nagasaki, Japan. *Tohoku J Exp Med* 164, 93-102, 1991.
- 6-200) Iwata K, Saito H, Nakano A. Association between cadmium-induced renal dysfunction and mortality: Further evidence. *Tohoku J Exp Med* 164, 319-330, 1991.
- 6-201) Arisawa K, Nakano A, Saito H, Liu X-J, Yokoo M, Soda M, Koba T, Takahashi T, Kinoshita K. Mortality and cancer incidence among a population previously exposed to environmental cadmium. *Int Arch Occup Environ Health* 74, 255-262, 2001.
- 6-202) イタイイタイ病および慢性カドミウム中毒に関する研究班, カドミウムによる環境汚染地域住民健康影響調査の解析および結果報告書, 日本公衆衛生協会, 1979.
- 6-203) Kawada T, Shinmyo R R, Suzuki S. Urinary cadmium and N-acetyl-β-D-glucosaminidase excretion of inhabitants living in a cadmium-polluted area. *Int Arch Occup Environ Health* 63, 541-546, 1992.
- 6-204) Nakadaira H, Nishi S. Effects of low-dose cadmium exposure on biological examinations. *Sci Total Environ* 308, 49-62, 2003.
- 6-205) Cai SW, Yue L, Hu ZN, Zhong XZ, Ye ZL, Xu HD, Liu YR, Ji RD, Zhang WH, Zhang FY. Cadmium exposure and health effects among residents in an irrigation area with ore dressing wastewater. *Sci Total Environ*. 1990 Jan;90:67-73.

- 6-206) Cai S, Yue L, Shang Q, Nordberg G. Cadmium exposure among residents in an area contaminated by irrigation water in China. *Bull World Health Organ.* 1995;73(3):359-67.
- 6-207) Han C. An investigation of the effects of cadmium exposure on human health. *Biomed Environ Sci.* 1988 Oct;1(3):323-31.
- 6-208) Jin T, Nordberg G, Wu X, Ye T, Kong Q, Wang Z, Zhuang F, Cai S. Urinary N-acetyl-beta-D-glucosaminidase isoenzymes as biomarker of renal dysfunction caused by cadmium in a general population. *Environ Res.* 1999 Aug;81(2):167-73.
- 6-209) Nordberg GF, Jin T, Kong Q, Ye T, Cai S, Wang Z, Zhuang F, Wu X. Biological monitoring of cadmium exposure and renal effects in a population group residing in a polluted area in China. *Sci Total Environ.* 1997 Jun 20;199(1-2):111-4.
- 6-210) Qu JB, Xin XF, Li SX, Ikeda M. Blood lead and cadmium in a general population in Jinan City, China. *Int Arch Occup Environ Health.* 1993;65(1 Suppl):S201-4.
- 6-211) Zhang ZW, Moon CS, Watanabe T, Shimbo S, He FS, Wu YQ, Zhou SF, Su DM, Qu JB, Ikeda M. Background exposure of urban populations to lead and cadmium: comparison between China and Japan. *Int Arch Occup Environ Health.* 1997;69(4):273-81.
- 6-212) Zhang ZW, Qu JB, Watanabe T, Shimbo S, Moon CS, Ikeda M. Exposure of citizens in China and in Japan to lead and cadmium: a comparative study. *Toxicol Lett.* 1999 Sep;108(2-3):167-72.

7. ヒトの健康影響評価

- 7-1) ATSDR.1999.Toxicological Profile for Cadmium. Atlanta GA: Agency for Toxic Substances and Disease Registry
- 7-2) Jarup L (ed) : Health effects of cadmium exposure – a review of the literature and a risk estimate. *Sacand J Work Environ Health* (1998) 24; suppl 1:52
- 7-3) Ikeda M, Ezaki T, Tsukahara T, Moriguchi J: Dietary cadmium intake in polluted and non-polluted areas in Japan in the past and in the present. *Int Arch Occup Environ Health* (2004), 77(4): 227-34
- 7-4) IPCS、Environmental Health Criteria 134 Cadmium (1992) p.22
- 7-5) IPCS、Environmental Health Criteria 134 Cadmium (1992) p.197
- 7-6) Grasseschi RM, Ramaswamy RB, Levine DJ, Klaassen CD, Wesselius LJ. Cadmium Accumulation and Detoxification by Alveolar Macrophages of Cigarette Smokers. *Chest* (2003), 124:1924-1928
- 7-7) Diamong GL, Thayer WC, Choudhury H: Pharmacokinetics/pharmacodynamics (PK/PD) modeling of risks of kidney toxicity from exposure to cadmium: estimates of dietary risks in the U.S. population. *J Toxicol Environ Health A*(2003), 66(22):2141-64
- 7-8) Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Dietary exposure to cadmium at close to the current provisional tolerable weekly intake does not affect renal function among female Japanese farmers. *Environmental Research* (2004) 95: 20-31
- 7-9) 水産物に含まれるカドミウムの実態調査結果について 平成15年5月2日水産庁

- 7-10) Kikuchi Y, Nomiyama T, Kumagai N, Uemura , Omae K. Cadmium Concentration in Current Japanese Food and Beverages. *J Occupa Health*(2002) 44;240247
- 7-11) Olsson IM, Bensryd I, Lundh T, Ottosson H, Skerfving S, Oskarsson A: Cadmium in Blood and Urine – Impact of Sex , Age, Dietary Intak, Iron Status, and Former Smoking – Association of Renal Effects. *Environ Health Persp* (2002), 110(12): 1185-1190
- 7-12) Watanabe T, Zhang ZW, Moon CS, Shimbo S Nakatsuka H, Matsuda-Inoguchi N, Higashikawa K, Ikeda M. Cadmium exposure of women in general populations in Japan during 1991-1997 compared with 1977-1981. *Int Arch Occup Environ Health*(2000) 73(1):26-34
- 7-13) Jung MC, Thomton I. Environmental contamination and seasonal variation of metals in soils, plants and waters in the paddy fields around a Pb-Zn mine in Korea. *Sci Total Environ*(1997) 198(2): 105-21
- 7-14) Sattrug S, Moore MR, Williams DJ. Cadmium Levels in the Lung, Liver, Kidney ,Cortes, and Urine Samples from Australians without Occupational Exposure to Metals. *Arch Environ Health* (2002) 57: 69-77
- 7-15) <http://minerals.usgs.gov/minerals/pubs/commodity/cadmium/index.html>
- 7-16) Ccortona G, Apostoli P, Toffoletto F, Baldasseroni A, Ghezzi I, Goggi E, Fornari S, Alessio L. Occupational exposure to cadmium and lung function. *Cadmium in the Human Environment: Toxicity and Carcinogenicity*, IARC 1992
- 7-17) Kawada T,Shinmyo RR,Suzuki S. Changes in unirmary cadmium excretion among pigment workers with improvement of the work environment. *Ind Health*(1993) 31:165-70
- 7-18) Wittman R, Hi H. Cadmium Exposure and Nephropathy in a 28-Year –Old Female Metals Worker. *Environ Health Persp* (2002) 110(12):1261-1266
- 7-19) http://www.jaish.or.jp/mms_s_shsi/13001
- 7-20) Fortoul TI, Osorio LS, Tovar AT, Salazar D, Castilla ME. Metal in Lung Tissue from Autopsy Cases in Mexico City Residents: Comparison Cases from the 1950s and the 1980s. *Environ Health Persp* (1996) 104(6): 630-632
- 7-21) Satarug S, Haswell-Elkins MR, Moore MR. Safe levels of cadmium intake to prevent renal tosicity in human subjects. *Br J of Nutr* (2000) 84:791-802
- 7-22) Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Comprehensive study of the effects of age, iron deficiency, diabetes mellitus, and cadmium burden on dietary cadmium absorption in cadmium-esposed female Japanese farmers. *Toxicology and Applied Pharmacology* (2004) 196: 114-123
- 7-23) Brzoska MM, Kaminski M, Dziki M, Moniuszko-Jakoniuk J. Changes in the structure and function of the kidney of rats chronically exposed to cadmium. II. Histoenzymatic studies. *Arch Toxicol*. 2004;78(4):226-31.
- 7-24) Brzoska MM, Kaminski M, Supernak-Bobko D, Zwierz K, Moniuszko-Jakoniuk J. Changes in the structure and function of the kidney of rats chronically exposed to cadmium. I. Biochemical and histopathological studies. *Arch Toxicol*. 2003;77(6):344-52.
- 7-25) Goyer RA, Clarkson TW. Toxic effects of metals (in Casarett and Doull's Toxicology, The Basic Science of Poison, 6th ed, ed Klaassen CD), p.811-67, McGraw-Hill, New York, 2001.
- 7-26) Honda R, Tsuritani I, Noborisaka Y, Suzuki H, Ishizaki M, Yamada Y. Urinary cadmium

- excretion is correlated with calcaneal bone mass in Japanese women living in an urban area. *Environ Res.* 2003;91(2):63-70.
- 7-27) Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Dietary exposure to cadmium at close to the current provisional tolerable weekly intake does not affect renal function among female Japanese farmers. *Environ Res.* 2004;95(1):20-31.
- 7-28) Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Environmental exposure to cadmium at a level insufficient to induce renal tubular dysfunction does not affect bone density among female Japanese farmers. *Environ Res.* in press.
- 7-29) Houtman JP. Prolonged low-level cadmium intake and atherosclerosis. *Sci Total Environ.* 1993;138(1-3):31-6.
- 7-30) Järup L, Berglund M, Elinder CG, Nordberg G, Vahter M. Health effects of cadmium exposure: a review of the literature and a risk estimate. *Scand J Work Environ Health.* 1998;24(Suppl 1):1-51.
- 7-31) 海津嘉蔵. イタイイタイ病の腎病変に関する調査研究. 環境保健レポート. 2000; 67:224-9.
- 7-32) 鍛冶利幸, 小山洋, 佐藤雅彦, 遠山千春. 低用量カドミウム暴露と健康影響 (2) 生活習慣病と生殖毒性. *日本衛生学雑誌.* 2002;57(3):556-63.
- 7-33) 小山洋, 鬼頭英明, 佐藤雅彦, 遠山千春. 低用量カドミウム暴露と健康影響 (1) 遺伝子傷害性と発がん性. *日本衛生学雑誌.* 2002;57(3):547-55.
- 7-34) 小山洋, 佐藤雅彦, 遠山千春. 低用量カドミウム暴露と健康影響 (4) 生体負荷量・臨界濃度・生命予後. *日本衛生学雑誌.* 2003;57(4):624-35.
- 7-35) Lyon TD, Aughey E, Scott R, Fell GS. Cadmium concentrations in human kidney in the UK: 1978-1993. *J Environ Monit.* 1999;1(3):227-31.
- 7-36) Nishijo M, Nakagawa H, Morikawa Y, Tabata M, Senma M, Miura K, Takahara H, Kawano S, Nishi M, Mizukoshi K, et al. Mortality of inhabitants in an area polluted by cadmium: 15 year follow up. *Occup Environ Med.* 1995;52(3):181-4.
- 7-37) Nogawa K, Honda R, Kido T, Tsuritani I, Yamada Y, Ishizaki M, Yamaya H. A dose-response analysis of cadmium in the general environment with special reference to total cadmium intake limit. *Environ Res.* 1989;48(1):7-16.
- 7-38) Satarug S, Baker JR, Urbenjapol S, Haswell-Elkins M, Reilly PE, Williams DJ, Moore MR. A global perspective on cadmium pollution and toxicity in non-occupationally exposed population. *Toxicol Lett.* 2003;137(1-2):65-83.
- 7-39) Sorahan T, Lancashire RJ. Lung cancer mortality in a cohort of workers employed at a cadmium recovery plant in the United States: an analysis with detailed job histories. *Occup Environ Med.* 1997;54(3):194-201.
- 7-40) Thevenod F. Nephrotoxicity and the proximal tubule. Insights from cadmium. *Nephron Physiol.* 2003;93(4):87-93.
- 7-41) Thun MJ, Osorio AM, Schober S, Hannon WH, Lewis B, Halperin W. Nephropathy in cadmium workers: assessment of risk from airborne occupational exposure to cadmium. *Br J Ind Med.* 1989;46(10):689-97.
- 7-42) Verougstraete V, Lison D, Hotz P. Cadmium, lung and prostate cancer: a systematic review of

- recent epidemiological data. *J Toxicol Environ Health B Crit Rev.* 2003;6(3):227-55.
- 7-43) WHO. Cadmium. Environmental Health Criteria 134. WHO, Geneva, 1992.
- 7-44) Elinder CG, Edling C, Lindberg E, Kagedal B, Vesterberg O. beta 2-Microglobulinuria among workers previously exposed to cadmium: follow-up and dose-response analyses. *Am J Ind Med.* 1985;8(6):553-64.
- 7-45) Holden H. Occupational Exposure of Cadmium. Cadmium Association. London. 1980:330-7.
- 7-46) Kjellstrom T. Accumulation and renal effects of cadmium in man. A dose-response study, Doctoral thesis, Department of Environmental Hygiene, Karolinska Institute. Stockholm, 1977a
- 7-47) Kjellstrom T, Evrin PE, Rahnster B. Dose-response analysis of cadmium-induced tubular proteinuria: a study of urinary beta2-microglobulin excretion among workers in a battery factory. *Environ Res.* 1977b Apr;13(2):303-17. No abstract available.
- 7-48) Roels H, Bernard AM, Cardenas A, Buchet JP, Lauwerys RR, Hotter G, Ramis I, Mutti A, Franchini I, Bundschuh I, et al. Markers of early renal changes induced by industrial pollutants. III. Application to workers exposed to cadmium. *Br J Ind Med.* 1993 Jan;50(1):37-48.
- 7-49) Toda K, Mori Y, Koike S, Ohmori K, Yamaga S. Chronic renal dysfunction in workers exposed to cadmium oxide fume. Medical examinations of welders using cadmium containing silver solder. *Sangyo Igaku.* 1984 May;26(3):212-23.
- 7-50) Thun MJ, Osorio AM, Schober S, Hannon WH, Lewis B, Halperin W. Nephropathy in cadmium workers: assessment of risk from airborne occupational exposure to cadmium. *Br J Ind Med.* 1989 Oct;46(10):689-97.
- 7-51) Alfvén, T., Järup, L., Elinder, C.G., 2002. Cadmium and lead in blood in relation to low bone mineral density and tubular proteinuria. *Environ. Health Perspect.* 110, 699–702.
- 7-52) Armitage, P., Berry, G., 1994. Statistical inference. In: *Statistical Methods in Medical Research*, 3rd Edition. Blackwell Science, Oxford, pp. 93–153.
- 7-53) Buchet, J.P., Lauwerys, R., Roels, H., Bernard, A., Bruaux, P., Claeys, F., Ducoffre, G., DePlaen, P., Staessen, J., Amery, A., Lijnen, P., Thijs, L., Rondia, D., Sartor, F., SaintRemy, A., Nick, L., 1990. Renal effects of cadmium body burden of the general population. *Lancet* 336, 699–702.
- 7-54) Hotz, P., Buchet, J.P., Bernard, A., Lison, D., Lauwerys, R., 1999. Renal effects of low-level environmental cadmium exposure: 5-year follow-up of a subcohort from the Cadmibel study. *Lancet* 354, 1508–1513.
- 7-55) Hotz, P., Buchet, J.P., Bernard, A., Lison, D., Lauwerys, R., 1999. Renal effects of low-level z environmental cadmium exposure: 5-year follow-up of a subcohort from the Cadmibel study. *Lancet* 354, 1508–1513.