

コホート研究・クロスセクショナル研究に係る参考文献一覧

番号	参考文献
1	Bellinger DC, Needleman HL, Leviton A, Waternaux C, Rabinowitz MB, Nichols ML. Early sensory-motor development and prenatal exposure to lead. Neurobehav Toxicol Teratol. 1984 Sep-Oct;6(5):387-402.
2	Bellinger DC, Leviton A, Needleman HL, Waternaux C, Rabinowitz M. Low-level lead exposure and infant development in the first year. Neurobehav Toxicol Tetatol.1986 Mar-Apr;8(2):151-61.
3	Bellinger DC, Leviton A, Waternaux C, Needleman H Rabinowitz M. Longitudinal analyses of prenatal and postnatal lead exposure and early cognitive development. N Engl J Med. 1987 Apr 23;316(17):1037-43.
4	Bellinger DC, Sloman J, Leviton A, Rabinowitz M, Needleman HL WaternauX C. Low-level lead exposure and children's cognitive function in the preschool years. Pediatrics. 1991 Feb; 87(2):219-27.
5	Bellinger DC, Stiles KM, Needleman HL. Low-level lead exposure, intelligence and academic achievement: a long-term follow-up study. Pediatrics. 1992 Dec;90(6):855-61.
6	Stiles KM, Bellinger DC. Neuropsychological correlates of low-level lead exposure in school-age children: a prospective study. Neurotoxicol Teratol. 1993 Jan-Feb;15(1):27-35.
7	Canfield RL, Henderson CR, Cory-Slechta DA, et al. Intellectual impairment in children with blood lead concentrations below 10 microgram per deciliter. N Engl J Med 348;16, 2003 Apr;17:1517-26.
8	Jusko TA., Henderson CR., Lanphear BP. et al. Blood lead concentrations <10 μg/dL and child intelligence at 6 Years of Age. Environ Health Perspect 2008 Feb;116(2)243-8.
9	Tellez-Rojo MM, Bellinger DC et al. Longitudinal associations between blood lead concentrations lower than 10 μg/dL and neurobehavioral development in environmentally exposed children in Mexico City. Pediatrics 2006 Aug;;118(2):e323-30.
10	Hu H, Tellez-Rojo MM, Bellinger D, et al. Fetal lead exposure at each stage of pregnancy as a predictor of infant mental development. Environ Health Perspect 2006 Nov;114(11):1730-35.
11	Schnass L et al. Reduced intellectual development in children with prenatal lead exposure. Environ Health Perspect 2006 May;114(5):791-97.
12	Surkan PJ, Zhang A, Trachtenberg F, Daniel DB, Mckinlay S, Bellinger DC, Neropsychological function in children with blood lead levels < 10 μg/dL. Neuro Toxicology 28, 2007;1170-1177.
13	Nigg JT, et al. Low blood lead levels associated with clinically diagnosed attention-deficit/hyperactivity disorder and mediated by weak cognitive control. BIOL PSYCHIATRY 2008;63:325-31.