

- B) the gene product(s)' function;
  - C) the phenotypic description of the new trait(s);
  - D) the level and site of expression in the plant of the expressed gene product(s), and the levels of its metabolites in the plant, particularly in the edible portions; and
  - E) where possible, the amount of the target gene product(s) if the function of the expressed sequence(s)/gene(s) is to alter the accumulation of a specific endogenous mRNA or protein.
33. In addition, information should be provided:
- A) to demonstrate whether the arrangement of the genetic material used for insertion has been conserved or whether significant rearrangements have occurred upon integration;
  - B) to demonstrate whether deliberate modifications made to the amino acid sequence of the expressed protein result in changes in its post-translational modification or affect sites critical for its structure or function;
  - C) to demonstrate whether the intended effect of the modification has been achieved and that all expressed traits are expressed and inherited in a manner that is stable through several generations consistent with laws of inheritance. It may be necessary to examine the inheritance of the DNA insert itself or the expression of the corresponding RNA if the phenotypic characteristics cannot be measured directly;
  - D) to demonstrate whether the newly expressed trait(s) are expressed as expected in the appropriate tissues in a manner and at levels that are consistent with the associated regulatory sequences driving the expression of the corresponding gene;
  - E) to indicate whether there is any evidence to suggest that one or several genes in the host plant has been affected by the transformation process; and
  - F) to confirm the identity and expression pattern of any new fusion proteins.

#### **SAFETY ASSESSMENT**

##### ***Expressed Substances (non-nucleic acid substances)***

##### *Assessment of possible toxicity*

34. *In vitro* nucleic acid techniques enable the introduction of DNA that can result in the synthesis of new substances in plants. The new substances can be conventional components of plant foods such as proteins, fats, carbohydrates, vitamins which are novel in the context of that recombinant-DNA plant. New substances might also include new metabolites resulting from the activity of enzymes generated by the expression of the introduced DNA.
35. The safety assessment should take into account the chemical nature and function of the newly expressed substance and identify the concentration of the substance in the edible parts of the recombinant-DNA plant, including variations and mean values. Current dietary exposure and possible effects on population sub-groups should also be considered.
36. Information should be provided to ensure that genes coding for known toxins or anti-nutrients present in the donor organisms are not transferred to recombinant-DNA plants that do not normally express those toxic or anti-nutritious characteristics. This assurance is particularly important in cases where a recombinant-DNA plant is processed differently from a donor plant, since conventional food processing techniques associated with the donor organisms may deactivate, degrade or eliminate anti-nutrients or toxicants.
37. For the reasons described in Section 3, conventional toxicology studies may not be considered necessary where the substance or a closely related substance has, taking into account its function and exposure,

been consumed safely in food. In other cases, the use of appropriate conventional toxicology or other studies on the new substance may be necessary.

38. In the case of proteins, the assessment of potential toxicity should focus on amino acid sequence similarity between the protein and known protein toxins and anti-nutrients (e.g. protease inhibitors, lectins) as well as stability to heat or processing and to degradation in appropriate representative gastric and intestinal model systems. Appropriate oral toxicity studies<sup>3</sup> may need to be carried out in cases where the protein present in the food is not similar to proteins that have previously been consumed safely in food, and taking into account its biological function in the plant where known.
39. Potential toxicity of non-protein substances that have not been safely consumed in food should be assessed on a case-by-case basis depending on the identity and biological function in the plant of the substance and dietary exposure. The type of studies to be performed may include studies on metabolism, toxicokinetics, sub-chronic toxicity, chronic toxicity/carcinogenicity, reproduction and development toxicity according to the traditional toxicological approach.
40. This may require the isolation of the new substance from the recombinant-DNA plant, or the synthesis or production of the substance from an alternative source, in which case, the material should be shown to be biochemically, structurally, and functionally equivalent to that produced in the recombinant-DNA plant.

#### *Assessment of possible allergenicity (proteins)*

41. When the protein(s) resulting from the inserted gene is present in the food, it should be assessed for potential allergenicity in all cases. An integrated, stepwise, case-by-case approach used in the assessment of the potential allergenicity of the newly-expressed protein(s) should rely upon various criteria used in combination (since no single criterion is sufficiently predictive on either allergenicity or non-allergenicity). As noted in paragraph 20, the data should be obtained using sound scientific methods. A detailed presentation of issues to be considered can be found in the Annex to this document.<sup>4</sup>
42. The newly expressed proteins in foods derived from recombinant-DNA plants should be evaluated for any possible role in the elicitation of gluten-sensitive enteropathy, if the introduced genetic material is obtained from wheat, rye, barley, oats, or related cereal grains.
43. The transfer of genes from commonly allergenic foods and from foods known to elicit gluten-sensitive enteropathy in sensitive individuals should be avoided unless it is documented that the transferred gene does not code for an allergen or for a protein involved in gluten-sensitive enteropathy.

#### *Compositional Analyses of Key Components*

44. Analyses of concentrations of key components<sup>5</sup> of the recombinant-DNA plant and, especially those typical of the food, should be compared with an equivalent analysis of a conventional counterpart grown and harvested under the same conditions. In some cases, a further comparison with the recombinant-DNA plant grown under its expected agronomic conditions may need to be considered (e.g. application of an herbicide). The statistical significance of any observed differences should be assessed in the context of the range of natural variations for that parameter to determine its biological significance. The comparator(s) used in this assessment should ideally be the near isogenic parental line. In practice, this

---

<sup>3</sup> Guidelines for oral toxicity studies have been developed in international fora, for example, the OECD Guidelines for the Testing of Chemicals.

<sup>4</sup> The FAO/WHO expert consultation 2001 report, which includes reference to several decision trees, was used in developing the Annex to these guidelines.

<sup>5</sup> Key nutrients or key anti-nutrients are those components in a particular food that may have a substantial impact in the overall diet. They may be major constituents (fats, proteins, carbohydrates as nutrients or enzyme inhibitors as anti-nutrients) or minor compounds (minerals, vitamins). Key toxicants are those toxicologically significant compounds known to be inherently present in the plant, such as those compounds whose toxic potency and level may be significant to health (e.g. solanine in potatoes if the level is increased, selenium in wheat) and allergens.

may not be feasible at all times, in which case a line as close as possible should be chosen. The purpose of this comparison, in conjunction with an exposure assessment as necessary, is to establish that substances that are nutritionally important or that can affect the safety of the food have not been altered in a manner that would have an adverse impact on human health.

45. The location of trial sites should be representative of the range of environmental conditions under which the plant varieties would be expected to be grown. The number of trial sites should be sufficient to allow accurate assessment of compositional characteristics over this range. Similarly, trials should be conducted over a sufficient number of generations to allow adequate exposure to the variety of conditions met in nature. To minimise environmental effects, and to reduce any effect from naturally occurring genotypic variation within a crop variety, each trial site should be replicated. An adequate number of plants should be sampled and the methods of analysis should be sufficiently sensitive and specific to detect variations in key components.

#### ***Evaluation of Metabolites***

46. Some recombinant-DNA plants may have been modified in a manner that could result in new or altered levels of various metabolites in the food. Consideration should be given to the potential for the accumulation of metabolites in the food that would adversely affect human health. Safety assessment of such plants requires investigation of residue and metabolite levels in the food and assessment of any alterations in nutrient profile. Where altered residue or metabolite levels are identified in foods, consideration should be given to the potential impacts on human health using conventional procedures for establishing the safety of such metabolites (e.g. procedures for assessing the human safety of chemicals in foods).

#### ***Food Processing***

47. The potential effects of food processing, including home preparation, on foods derived from recombinant-DNA plants should also be considered. For example, alterations could occur in the heat stability of an endogenous toxicant or the bioavailability of an important nutrient after processing. Information should therefore be provided describing the processing conditions used in the production of a food ingredient from the plant. For example, in the case of vegetable oil, information should be provided on the extraction process and any subsequent refining steps.

#### ***Nutritional Modification***

48. The assessment of possible compositional changes to key nutrients, which should be conducted for all recombinant-DNA plants, has already been addressed under 'Compositional analyses of key components'. However, foods derived from recombinant-DNA plants that have undergone modification to intentionally alter nutritional quality or functionality should be subjected to additional nutritional assessment to assess the consequences of the changes and whether the nutrient intakes are likely to be altered by the introduction of such foods into the food supply.
49. Information about the known patterns of use and consumption of a food, and its derivatives should be used to estimate the likely intake of the food derived from the recombinant-DNA plant. The expected intake of the food should be used to assess the nutritional implications of the altered nutrient profile both at customary and maximal levels of consumption. Basing the estimate on the highest likely consumption provides assurance that the potential for any undesirable nutritional effects will be detected. Attention should be paid to the particular physiological characteristics and metabolic requirements of specific population groups such as infants, children, pregnant and lactating women, the elderly and those with chronic diseases or compromised immune systems. Based on the analysis of nutritional impacts and the dietary needs of specific population subgroups, additional nutritional assessments may be necessary. It is also important to ascertain to what extent the modified nutrient is bioavailable and remains stable with time, processing and storage.
50. The use of plant breeding, including *in vitro* nucleic acid techniques, to change nutrient levels in crops can result in broad changes to the nutrient profile in two ways. The intended modification in plant constituents could change the overall nutrient profile of the plant product and this change could affect the nutritional status of individuals consuming the food. Unexpected alterations in nutrients could have the

same effect. Although the recombinant-DNA plant components may be individually assessed as safe, the impact of the change on the overall nutrient profile should be determined.

51. When the modification results in a food product, such as vegetable oil, with a composition that is significantly different from its conventional counterpart, it may be appropriate to use additional conventional foods or food components (i.e. foods or food components whose nutritional composition is closer to that of the food derived from recombinant-DNA plant) as appropriate comparators to assess the nutritional impact of the food.
52. Because of geographical and cultural variation in food consumption patterns, nutritional changes to a specific food may have a greater impact in some geographical areas or in some cultural population than in others. Some food plants serve as the major source of a particular nutrient in some populations. The nutrient and the populations affected should be identified.
53. Some foods may require additional testing. For example, animal feeding studies may be warranted for foods derived from recombinant-DNA plants if changes in the bioavailability of nutrients are expected or if the composition is not comparable to conventional foods. Also, foods designed for health benefits may require specific nutritional, toxicological or other appropriate studies. If the characterization of the food indicates that the available data are insufficient for a thorough safety assessment, properly designed animal studies could be requested on the whole foods.

### **SECTION 5 – OTHER CONSIDERATIONS**

#### **POTENTIAL ACCUMULATION OF SUBSTANCES SIGNIFICANT TO HUMAN HEALTH**

54. Some recombinant-DNA plants may exhibit traits (e.g., herbicide tolerance) which may indirectly result in the potential for accumulation of pesticide residues, altered metabolites of such residues, toxic metabolites, contaminants, or other substances which may be relevant to human health. The safety assessment should take this potential for accumulation into account. Conventional procedures for establishing the safety of such compounds (e.g., procedures for assessing the human safety of chemicals) should be applied.

#### **USE OF ANTIBIOTIC RESISTANCE MARKER GENES**

55. Alternative transformation technologies that do not result in antibiotic resistance marker genes in foods should be used in the future development of recombinant-DNA plants, where such technologies are available and demonstrated to be safe.
56. Gene transfer from plants and their food products to gut microorganisms or human cells is considered a rare possibility because of the many complex and unlikely events that would need to occur consecutively. Nevertheless, the possibility of such events cannot be completely discounted<sup>6</sup>.
57. In assessing safety of foods containing antibiotic resistance marker genes, the following factors should be considered:
  - A) the clinical and veterinary use and importance of the antibiotic in question;  
(Certain antibiotics are the only drug available to treat some clinical conditions (e.g. vancomycin for use in treating certain staphylococcal infections). Marker genes encoding resistance to such antibiotics should not be used in recombinant-DNA plants.)
  - B) whether the presence in food of the enzyme or protein encoded by the antibiotic resistance marker gene would compromise the therapeutic efficacy of the orally administered antibiotic; and  
(This assessment should provide an estimate of the amount of orally ingested antibiotic that could be degraded by the presence of the enzyme in food, taking into account factors such as dosage of the antibiotic, amount of enzyme likely to remain in food following exposure to digestive conditions,

---

<sup>6</sup> In cases where there are high levels of naturally occurring bacteria which are resistant to the antibiotic, the likelihood of such bacteria transferring this resistance to other bacteria will be orders of magnitude higher than the likelihood of transfer between ingested foods and bacteria.

including neutral or alkaline stomach conditions and the need for enzyme cofactors (e.g. ATP) for enzymatic activity and estimated concentration of such factors in food.)

C) safety of the gene product, as would be the case for any other expressed gene product.

58. If evaluation of the data and information suggests that the presence of the antibiotic resistance marker gene or gene product presents risks to human health, the marker gene or gene product should not be present in the food. Antibiotic resistance genes used in food production that encode resistance to clinically used antibiotics should not be present in foods.

#### ***REVIEW OF SAFETY ASSESSMENTS***

59. The goal of the safety assessment is a conclusion as to whether the new food is as safe as the conventional counterpart taking into account dietary impact of any changes in nutritional content or value. Nevertheless, the safety assessment should be reviewed in the light of new scientific information that calls into question the conclusions of the original safety assessment.