

<出典>

英国 “The BSE Inquiry, Vol.16, p37, October 2000”

Figure 3.9: Number of confirmed cases of BSE by age at clinical onset, and by year of clinical onset, Great Britain

Year of clinical onset	Year of age at clinical onset															
	1*	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
1986	0	2	4	3	2	0	0	0	0	0	0	0	0	0	0	0
1987	0	4	78	171	87	14	11	0	1	0	0	0	0	0	0	0
1988	0	16	354	1,124	806	211	50	22	5	3	2	0	0	0	0	0
1989	1	31	661	2,356	2,150	782	146	40	10	2	3	0	0	0	1	0
1990	0	54	1,614	4,074	3,006	1,403	394	73	23	12	5	0	3	0	0	0
1991	0	27	3,110	8,060	4,680	1,807	647	169	65	14	9	2	1	1	0	0
1992	1	25	1,224	11,414	9,121	2,977	815	327	114	21	9	8	0	1	0	0
1993	0	17	841	5,290	10,980	5,306	1,270	388	157	67	22	3	2	0	0	0
1994	0	10	400	3,169	5,395	5,461	2,047	522	197	85	36	4	5	0	0	0
1995	0	5	413	1,725	3,284	2,881	1,919	704	213	76	32	17	4	1	0	0
1996	0	1	46	301	348	365	276	135	40	17	4	7	0	0	0	0

*That is, between 12 and 24 months.

Year of clinical onset	Total age	Year of age at clinical onset		Total
		Mean age (years)	age unknown	
1986	11	3.5	1	12
1987	366	4.2	94	460
1988	2,593	4.4	551	3,144
1989	6,183	4.6	1,592	7,775
1990	10,661	4.5	3,951	14,612
1991	18,592	4.4	7,264	25,856
1992	26,057	4.7	11,094	37,151
1993	24,343	5.1	9,428	33,771
1994	17,331	5.5	5,579	22,910
1995	11,274	5.7	2,497	13,771
1996	1,540	5.8	241	1,781
Total	161,243			

Results shown as percentages

Year of clinical onset	Year of age at clinical onset															
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
1986	0%	18%	36%	27%	18%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
1987	0%	1%	21%	47%	24%	4%	3%	0%	0%	0%	0%	0%	0%	0%	0%	0%
1988	0%	1%	14%	43%	31%	8%	2%	1%	0%	0%	0%	0%	0%	0%	0%	0%
1989	0%	1%	11%	38%	35%	13%	2%	1%	0%	0%	0%	0%	0%	0%	0%	0%
1990	0%	1%	15%	38%	28%	13%	4%	1%	0%	0%	0%	0%	0%	0%	0%	0%
1991	0%	0%	17%	43%	25%	10%	3%	1%	0%	0%	0%	0%	0%	0%	0%	0%
1992	0%	0%	5%	44%	35%	11%	3%	1%	0%	0%	0%	0%	0%	0%	0%	0%
1993	0%	0%	3%	22%	45%	22%	5%	2%	1%	0%	0%	0%	0%	0%	0%	0%
1994	0%	0%	2%	18%	31%	32%	12%	3%	1%	0%	0%	0%	0%	0%	0%	0%
1995	0%	0%	4%	15%	29%	26%	17%	6%	2%	1%	0%	0%	0%	0%	0%	0%
1996	0%	0%	3%	20%	23%	24%	18%	9%	3%	1%	0%	0%	0%	0%	0%	0%

Oldest and youngest cases

Although the number was small, there were a few cases in which clinical onset occurred between 20 and 30 months. The youngest was indeed just 20 months old (in England in 1992). The oldest case was 18 years old (in 1993).