

「8.2.3 疫学調査結果」に係る参考文献リスト

- No.1 Ikeda M, Ezaki T, Moriguchi J, Fukui Y, Ukai H, Okamoto S, Sakurai H
The Threshold Cadmium Level That Causes a Substantial Increase in β 2-Microglobulin
in Urine of General Populations
Tohoku J. Exp. Med., 2005,205, 247-261
- No.2 Ikeda M., Ezaki T., Tsukahara T., Moriguchi J., Furuki K., Fukui Y., Ukai H., Okamoto S., Sakurai H.
Threshold levels of urinary cadmium in relation to increases in urinary β 2-microglobulin
among general Japanese populations.
Toxicol. Lett. 2003;137:135-141.
- No.3 Gamo M, Ono K, Nakanishi J.
Meta-analysis for deriving age- and gender-specific dose-response relationships between
urinary cadmium concentration and β 2-microglobulinuria under environmental exposure.
Environmental Research 101 (2006) 104-112.
- No.4 Nogawa K, Honda R, Kido T, Tsuritani I, Yamada Y, Ishizaki M, Yamaya H.
A Dose-Response Analysis of Cadmium in the General Environment with Special Reference
to Total Cadmium Intake Limit.
Environ Res. 1989; 48, 7-16
- No.5 Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F.
Dietary exposure to cadmium at close to the current provisional tolerable weekly intake
dose not affect renal function among female Japanese farmers.
Environ Res. 2004 May; 95 (1): 20-31.