## 選定された文献(疫学)

| 通し<br>No. | エンドポイント | 評価値算出機関  | 区分                       | タイトル  | 年     | 著者                               | 雑誌                       |
|-----------|---------|--|--------------------------|---|-------|----------------------------------|--------------------------|
| 1         | 免疫      | EFSA 2020  | POD                      | Internal exposure to perfluoroalkyl substances (PFASs) and biological markers in 101 healthy 1-year-old children: associations between levels of perfluorooctanoic acid (PFOA) and vaccine response             | 2020  | Abraham et al.                   | Arch Toxicol             |
| 2         | 免疫      | EPA 2021<br>(PFOS,<br>PFOA)<br>EPA 2023<br>(PFOS,<br>PFOA) | POD                      | Application of benchmark analysis for mixed contaminant exposures: Mutual adjustment of perfluoroalkylate substances associated with immunotoxicity   | 2018  | Budtz-Jørgensen<br>and Grandjean | PLoS One                 |
| 3         | 免疫      | EPA 2023<br>(PFOS,<br>PFOA)                                | POD候<br>補                | Concentrations of tetanus and diphtheria antibodies in vaccinated Greenlandic children aged 7-12 years exposed to marine pollutants, a cross sectional study.   | 2021  | Timmerman et al.                 | Environ Res              |
| 4         | 免疫      | EPA 2021<br>(PFOS,<br>PFOA)                                | 過去の<br>POD               | Estimated exposures to perfluorinated compounds in infancy predict attenuated vaccine antibody concentrations at age 5-years  | 2017a | Grandjean et al.                 | J Immunotoxicol          |
| 5         | 免疫      | EPA 2021<br>(PFOS,<br>PFOA)                                | 過去の<br>POD               | Serum vaccine antibody concentrations in adolescents exposed to perfluorinated compounds  | 2017b | Grandjean et al.                 | Environ Health Perspect  |
| 6         | 免疫      | EPA 2021<br>(PFOS,<br>PFOA)<br>EFSA 2018                   | 過去の<br>POD               | Serum vaccine antibody concentrations in children exposed to perfluorinated compounds   | 2012  | Grandjean et al.                 | JAMA                     |
| 7         | 免疫      |  | 専門委<br>員/専<br>門参考<br>人選択 | Prenatal exposure to perfluoroalkyl and polyfluoroalkyl substances and childhood atopic dermatitis: a prospective birth cohort study  | 2018  | Chen et al.                      | Environ Health           |
| 8         | 免疫      |  |                          | Effect of prenatal exposure to per- and polyfluoroalkyl substances on childhood allergies and common infectious diseases in children up to age 7 years: The Hokkaido study on environment and children's health | 2020  | Ait Bamai et al.                 | Environ Int              |
| 9         | 免疫      |  |                          | Serum polyfluoroalkyl concentrations, asthma outcomes, and immunological markers in a case-control study of Taiwanese children  | 2013  | Dong et al.                      | Environ Health Perspect  |
| 10        | 免疫      |  |                          | PFAS (per- and polyfluoroalkyl substances) and asthma in young children: NHANES 2013-2014   | 2020  | Jackson-Browne<br>et al.         | Int J Hyg Environ Health |
| 11        | 免疫      |  |                          | Exposure to perfluoroalkyl substances and allergic outcomes in children: A systematic review and meta-analysis  | 2020  | Luo et al.                       | Environ Res              |
| 12        | 免疫      |  |                          | Exposure to perfluoroalkyl substances and neonatal immunoglobulin profiles in the upstate KIDS study (2008-2010)  | 2022  | Jones et al.                     | Environ Pollut           |
| 13        | 免疫      |  |                          | Effects of exposure to per- and polyfluoroalkyl substances on vaccine antibodies: A systematic review and meta-analysis based on epidemiological studies  | 2022  | Zhang et al.                     | Environ Pollut           |

| 通し<br>No. | エンドポイント | 評価値算出 機関 | 区分 | タイトル   | 年    | 著者                        | 雑誌                       |
|-----------|---------|----------|----|--|------|---------------------------|--------------------------|
| 14        | 免疫      |          |    | Prenatal exposure to perfluorinated chemicals and relationship with allergies and infectious diseases in infants   | 2012 | Okada et al.              | Environ Res              |
| 15        | 免疫      |          |    | Prenatal exposure to perfluoroalkyl acids and prevalence of infectious diseases up to 4 years of age   | 2017 | Goudarzi et al.           | Environ Int              |
| 16        | 免疫      |          |    | Maternal levels of perfluoroalkyl substances (PFASs) during pregnancy and childhood allergy and asthma related outcomes and infections in the Norwegian Mother and Child (MoBa) cohort | 2019 | Impinen et al.            | Environ Int              |
| 17        | 免疫      |          |    | Prenatal exposure to perfluoralkyl substances (PFASs) associated with respiratory tract infections but not allergy- and asthma-related health outcomes in childhood                    | 2018 | Impinen et al.            | Environ Res              |
| 18        | 免疫      |          |    | Prenatal exposure to perfluoroalkyl substances, immune-related outcomes, and lung function in children from a Spanish birth cohort study   | 2019 | Manzano-Salgado<br>et al. | Int J Hyg Environ Health |
| 19        | 免疫      |          |    | Prenatal perfluorooctanoic acid exposure is associated with early onset atopic dermatitis in 5-year-old children   | 2019 | Wen et al.                | Chemosphere              |
| 20        | 免疫      |          |    | Exposure to perfluoroalkyl substances during fetal life and hospitalization for infectious disease in childhood: A study among 1,503 children from the Odense Child Cohort.            | 2021 | Dalsager et al.           | Environ Int              |
| 21        | 免疫      |          |    | Cord blood gene expression supports that prenatal exposure to perfluoroalkyl substances causes depressed immune functionality in early childhood                                       | 2016 | Pennings et al.           | J Immunotoxicol          |

| 通し<br>No. | エンドポイント   | 評価値算出機関                     | 区分                       | タイトル  | 年    | 著者                 | 雑誌                              |
|-----------|-----------|-----------------------------|--------------------------|---|------|--------------------|---------------------------------|
| 22        | 生殖・発<br>生 | EPA 2023<br>(PFOS,<br>PFOA) | POD                      | Maternal serum levels of perfluoroalkyl substances in early pregnancy and offspring birth weight.   | 2020 | Wikström et al.    | Pediatr Res                     |
| 23        | 生殖・発<br>生 | EPA 2023<br>(PFOS,<br>PFOA) | POD候<br>補                | Early-Pregnancy Plasma Concentrations of<br>Perfluoroalkyl Substances and Birth Outcomes in<br>Project Viva: Confounded by Pregnancy<br>Hemodynamics?   | 2018 | Sagiv et al.       | Am J Epidemiol                  |
| 24        | 生殖・発<br>生 | EPA 2023<br>(PFOS,<br>PFOA) | POD候<br>補                | Perfluoroalkyl substances during pregnancy and offspring weight and adiposity at birth: Examining mediation by maternal fasting glucose in the healthy start study  | 2017 | Starling et al.    | Environ Health Perspect         |
| 25        | 生殖・発<br>生 | EPA 2023<br>(PFOS,<br>PFOA) | POD候<br>補                | Are perfluorooctane sulfonate alternatives safer? New insights from a birth cohort study.   | 2020 | Chu et al.         | Environ Int                     |
| 26        | 生殖・発<br>生 | EPA 2023<br>(PFOS)          | POD候<br>補                | Serum perfluorooctanoic acid and perfluorooctane sulfonate concentrations in relation to birth outcomes in the Mid-Ohio Valley, 2005-2010.  | 2013 | Darrow et al.      | Environ Health Perspect         |
| 27        | 生殖・発<br>生 | EPA 2023<br>(PFOS)          | POD候<br>補                | Associations of paternal and maternal per- and polyfluoroalkyl substances exposure with cord serum reproductive hormones, placental steroidogenic enzyme and birth weight.                                | 2021 | Yao et al.         | Chemosphere                     |
| 28        | 生殖・発<br>生 | EPA 2023<br>(PFOA)          | POD候<br>補                | Combined Effects of Prenatal Exposures to Environmental Chemicals on Birth Weight.  | 2016 | Govarts et al.     | Int J Environ Res Public Health |
| 29        | 生殖・発<br>生 |                             | 専門委<br>員/専<br>門参考<br>人選択 | Prenatal exposure to perfluorooctanoate and risk of overweight at 20 years of age: a prospective cohort study   | 2012 | Halldorsson et al. | Environ Health Perspect         |
| 30        | 生殖・発<br>生 |                             |                          | Cumulative exposure to environmental pollutants<br>during early pregnancy and reduced fetal growth: the<br>Project Viva cohort  | 2018 | Rokoff et al.      | Environ Health                  |
| 31        | 生殖・発<br>生 |                             |                          | Prenatal exposure to per- and polyfluoroalkyl substances and infant growth and adiposity: The healthy start study   | 2019 | Starling et al.    | Environ Int                     |
| 32        | 生殖・発<br>生 |                             |                          | Associations of Perfluoroalkyl Substances (PFAS) with lower birth weight: an evaluation of potential confounding by glomerular filtration rate using a physiologically based pharmacokinetic model (PBPK) | 2015 | Verner et al.      | Environ Health Perspect         |
| 33        | 生殖・発<br>生 |                             |                          | Gestational perfluoroalkyl substance exposure and<br>body mass index trajectories over the first 12 years of<br>life  | 2021 | Braun et al.       | Int J Obes (Lond)               |
| 34        | 生殖・発<br>生 |                             |                          | Attention deficit/hyperactivity disorder and childhood autism in association with prenatal exposure to perfluoroalkyl substances: A nested case-control study in the Danish National Birth Cohort         | 2015 | Liew et al.        | Environ Health Perspect         |

| 通し<br>No. | エンドポイント   | 評価値算出機関 | 区分 | タイトル  | 年    | 著者                        | 雑誌                         |
|-----------|-----------|---------|----|---|------|---------------------------|----------------------------|
| 35        | 生殖・発<br>生 |         |    | Early prenatal exposure to suspected endocrine disruptor mixtures is associated with lower IQ at age seven  | 2020 | Tanner et al.             | Environ Int                |
| 36        | 生殖・発<br>生 |         |    | Concentrations of perfluoroalkyl substances and bisphenol A in newborn dried blood spots and the association with child behavior  | 2018 | Ghassabian et al.         | Environ Pollut             |
| 37        | 生殖・発<br>生 |         |    | Prenatal and childhood exposure to per- and polyfluoroalkyl substances (PFASs) and child cognition  | 2018 | Harris et al.             | Environ Int                |
| 38        | 生殖・発<br>生 |         |    | The association between prenatal exposure to perfluoroalkyl substances and childhood neurodevelopment   | 2020 | Spratlen et al.           | Environ Pollut             |
| 39        | 生殖・発<br>生 |         |    | Prenatal and childhood exposure to poly- and perfluoroalkyl substances (PFAS) and cognitive development in children at age 8 years  | 2019 | Vuong et al.              | Environ Res                |
| 40        | 生殖・発<br>生 |         |    | The association between prenatal endocrine-<br>disrupting chemical exposure and altered resting-<br>state brain fMRI in teenagers   | 2020 | Weng et al.               | Brain Struct Funct         |
| 41        | 生殖・発<br>生 |         |    | The association between maternal perfluoroalkyl substances exposure and early attention deficit hyperactivity disorder in children: a systematic review and meta-analysis   | 2021 | Qu et al.                 | Environ Sci Pollut Res Int |
| 42        | 生殖・発<br>生 |         |    | Early life multiple exposures and child cognitive function: A multi-centric birth cohort study in six European countries  | 2021 | Julvez et al.             | Environ Pollut             |
| 43        | 生殖・発<br>生 |         |    | The association between prenatal perfluoroalkyl substance exposure and symptoms of attention-deficit/hyperactivity disorder in 8-year-old children and the mediating role of thyroid hormones in the Hokkaido study | 2022 | Itoh et al.               | Environ Int                |
| 44        | 生殖・発<br>生 |         |    | Prenatal exposure to perfluoroalkyl substances and cardiometabolic risk in children from the Spanish INMA birth cohort study  | 2017 | Manzano-Salgado<br>et al. | Environ Health Perspect    |
| 45        | 生殖・発<br>生 |         |    | Gestational exposure to perfluoroalkyl substances and congenital heart defects: A nested case-control pilot study   | 2021 | Ou et al.                 | Environ Int                |
| 46        | 生殖・発<br>生 |         |    | Prenatal exposure to per- and polyfluoroalkyl substances and childhood adiposity at 7 years of age  | 2022 | Zhang et al.              | Chemosphere                |
| 47        | 生殖・発<br>生 |         |    | Prenatal Exposure to Perfluoroalkyl Substances<br>Associated With Increased Susceptibility to Liver<br>Injury in Children   | 2020 | Stratakis et al.          | Hepatology                 |
| 48        | 生殖・発<br>生 |         |    | Association of Prenatal Exposure to Endocrine-<br>Disrupting Chemicals With Liver Injury in Children  | 2022 | Midya et al.              | JAMA Netw Open             |

| 通し<br>No. | エンドポイント   | 評価値算出 機関 | 区分 | タイトル  | 年    | 著者                | 雑誌                      |
|-----------|-----------|----------|----|---|------|-------------------|-------------------------|
| 49        | 生殖・発<br>生 |          |    | Exposure to perfluoroalkyl substances during fetal life and pubertal development in boys and girls from the danish national birth cohort  | 2019 | Ernst et al.      | Environ Health Perspect |
| 50        | 生殖・発<br>生 |          |    | Cord blood per- and polyfluoroalkyl substances, placental steroidogenic enzyme, and cord blood reproductive hormone   | 2019 | Yao et al.        | Environ Int             |
| 51        | 生殖・発<br>生 |          |    | Early life exposures to perfluoroalkyl substances in relation to adipokine hormone levels at birth and during childhood   | 2019 | Shelly et al.     | J Clin Endocrinol Metab |
| 52        | 生殖・発<br>生 |          |    | Associations of perfluoroalkyl substances with adipocytokines in umbilical cord serum: A mixtures approach  | 2022 | Ding et al.       | Environ Res             |
| 53        | 生殖・発<br>生 |          |    | Exposure to Perflouroalkyl acids and foetal and maternal thyroid status: a review   | 2020 | Boesen et al.     | Environ Health          |
| 54        | 生殖・発<br>生 |          |    | Association of exposure to prenatal perfluoroalkyl substances and estrogen receptor 1 polymorphisms with the second to fourth digit ratio in school-aged children: The Hokkaido study   | 2022 | Nishimura et al.  | Reprod Toxicol          |
| 55        | 生殖・発<br>生 |          |    | Associations of per- and polyfluoroalkyl substances (PFAS) and their mixture with oxidative stress biomarkers during pregnancy  | 2022 | Taibl et al.      | Environ Int             |
| 56        | 生殖・発<br>生 |          |    | Prenatal exposure to mixtures of persistent endocrine disrupting chemicals and early menarche in a population-based cohort of British girls   | 2021 | Marks et al.      | Environ Pollut          |
| 57        | 生殖・発<br>生 |          |    | Endocrine disruptors and neonatal anthropometry,<br>NICHD Fetal Growth Studies - Singletons   | 2018 | Buck Louis et al. | Environ Int             |
| 58        | 生殖・発<br>生 |          |    | Prenatal Exposure to Per- and Polyfluoroalkyl Substances (PFASs) and Association between the Placental Transfer Efficiencies and Dissociation Constant of Serum Proteins-PFAS Complexes | 2019 | Gao et al.        | Environ Sci Technol     |
| 59        | 生殖・発<br>生 |          |    | Exposure to perfluorinated compounds and human semen quality in arctic and European populations   | 2012 | Toft et al.       | Hum Reprod              |
| 60        | 生殖・発<br>生 |          |    | Association of perfluoroalkyl and polyfluoroalkyl substances with premature ovarian insufficiency in Chinese women  | 2018 | Zhang et al.      | J Clin Endocrinol Metab |
| 61        | 生殖・発<br>生 |          |    | The effects of perfluoroalkyl and polyfluoroalkyl substances on female fertility: A systematic review and meta-analysis   | 2022 | Wang et al.       | Environ Res             |
| 62        | 生殖・発<br>生 |          |    | Global Exposure to Per- and Polyfluoroalkyl<br>Substances and Associated Burden of Low<br>Birthweight   | 2022 | Fan et al.        | Environ Sci Technol     |

| 通し<br>No. | エンドポイント | 評価値算出 機関 | 区分 | タイトル  | 年    | 著者          | 雑誌                         |
|-----------|---------|----------|----|---|------|-------------|----------------------------|
| 6:        | 生殖・発生   |          |    | Per- and polyfluoroalkyl substances exposure during pregnancy and adverse pregnancy and birth outcomes: A systematic review and meta-analysis | 2021 | Gao et al.  | Environ Res                |
| 64        | 生殖・発生   |          |    | Association between per- and polyfluoroalkyl substances and semen quality   | 2022 | Wang et al. | Environ Sci Pollut Res Int |
| 6         | 生殖・発生   |          |    | Associations between exposure to perfluoroalkyl substances and birth outcomes: A meta-analysis  | 2022 | Yang et al. | Chemosphere                |

| 通し<br>No. | エンドポイント | 評価値算出機関                                  | 区分                       | タイトル  | 年    | 著者                          | 雑誌                      |
|-----------|---------|--|--------------------------|---|------|-----------------------------|-------------------------|
| 66        | 代謝      | EPA 2023<br>(PFOS,<br>PFOA)              | POD                      | Using 2003-2014 U.S. NHANES data to determine the associations between per- and polyfluoroalkyl substances and cholesterol: Trend and implications.   | 2019 | Dong et al.                 | Ecotoxicol Environ Saf  |
| 67        | 代謝      | EPA 2023<br>(PFOS,<br>PFOA)<br>EFSA 2018 | POD候<br>補                | Association of perfluorooctanoic acid and perfluorooctane sulfonate with serum lipids among adults living near a chemical plant   | 2009 | Steenland et al.            | Am J Epidemiol          |
| 68        | 代謝      | EPA 2023<br>(PFOS,<br>PFOA)              | POD候<br>補                | Per- and polyfluoroalkyl substances and blood lipid levels in pre-diabetic adults-longitudinal analysis of the diabetes prevention program outcomes study.                                      | 2019 | Lin et al.                  | Environ Int             |
| 69        | 代謝      | EFSA 2018                                | 過去の<br>POD               | Association between plasma PFOA and PFOS levels and total cholesterol in a middle-aged Danish population  | 2013 | Eriksen et al.              | PLoS One                |
| 70        | 代謝      | EFSA 2018                                | 過去の<br>POD               | Exposure to Polyfluoroalkyl Chemicals and<br>Cholesterol, Body Weight, and Insulin Resistance in<br>the General US Population   | 2010 | Nelson et al.               | Environ Health Perspect |
| 71        | 代謝      |  | 専門委<br>員/専<br>門参考<br>人選択 | Stochastic pharmacokinetic-pharmacodynamic modeling for assessing the systemic health risk of perfluorooctanoate (pfoa)   | 2018 | Convertino et al.           | Toxicol Sci             |
| 72        | 代謝      |  |                          | The association between PFOA, PFOS and serum lipid levels in adolescents  | 2021 | Geiger et al.               | Chemosphere             |
| 73        | 代謝      |  |                          | The association between perfluoroalkyl chemicals and serum lipid levels in children   | 2015 | Zeng et al.                 | Sci Total Environ       |
| 74        | 代謝      |  |                          | Serum albumin mediates the effect of multiple per-<br>and polyfluoroalkyl substances on serum lipid levels  | 2020 | Fan et al.                  | Environ Pollut          |
| 75        | 代謝      |  |                          | Associations between lipid/lipoprotein levels and perfluoroalkyl substances among US children aged 6-11 years   | 2018 | Jain et al.                 | Environ Pollut          |
| 76        | 代謝      |  |                          | The association between total serum isomers of per-<br>and polyfluoroalkyl substances, lipid profiles, and the<br>DNA oxidative/nitrative stress biomarkers in middle-<br>aged Taiwanese adults | 2020 | Lin et al.                  | Environ Res             |
| 77        | 代謝      |  |                          | Association among total serum isomers of perfluorinated chemicals, glucose homeostasis, lipid profiles, serum protein and metabolic syndrome in adults: NHANES, 2013-2014                       | 2018 | Liu et al.                  | Environ Pollut          |
| 78        | 代謝      |  |                          | Exposure to Perfluoroalkyl Substances and Metabolic<br>Outcomes in Pregnant Women: Evidence from the<br>Spanish INMA Birth Cohorts  | 2017 | Matilla-Santander<br>et al. | Environ Health Perspect |
| 79        | 代謝      |  |                          | Examining confounding by diet in the association between perfluoroalkyl acids and serum cholesterol in pregnancy  | 2015 | Skuladottir et al.          | Environ Res             |

| 通し<br>No. | エンドポイ<br>ント | 評価値算出 機関                    | 区分        | タイトル  | 年    | 著者                 | 雑誌                              |
|-----------|-------------|-----------------------------|-----------|---|------|--------------------|---------------------------------|
| 80        | 代謝          |                             |           | Perfluoroalkyl substance mixtures and cardio-<br>metabolic outcomes in highly exposed male workers<br>in the Veneto Region: A mixture-based approach  | 2022 | Batzella et al.    | Environ Res                     |
| 81        | 代謝          |                             |           | PFAS Concentrations and Cardiometabolic Traits in<br>Highly Exposed Children and Adolescents  | 2021 | Canova et al.      | Int J Environ Res Public Health |
| 82        | 代謝          |                             |           | The PFAS Health Study: Systematic Literature Review   | 2018 | Kirk et al.        | -                               |
| 83        | 代謝          |                             |           | Adiposity and glycemic control in children exposed to perfluorinated compounds  | 2014 | Timmermann et al.  | J Clin Endocrinol Metab         |
| 84        | 代謝          |                             |           | Perfluoroalkyl substances, metabolomic profiling, and alterations in glucose homeostasis among overweight and obese Hispanic children: A proof-of-concept analysis  | 2019 | Alderete et al.    | Environ Int                     |
| 85        | 代謝          |                             |           | Longitudinal associations of exposure to perfluoroalkylated substances in childhood and adolescence and indicators of adiposity and glucose metabolism 6 and 12 years later: The European Youth Heart Study | 2016 | Domazet et al.     | Diabetes Care                   |
| 86        | 代謝          |                             |           | The association between perfluoroalkyl substances and lipids in cord blood  | 2020 | Spratlen et al.    | J Clin Endocrinol Metab         |
| 87        | 代謝          |                             |           | Environmental exposure to perfluoroalkyl substances in early pregnancy, maternal glucose homeostasis and the risk of gestational diabetes: A prospective cohort study                                       | 2021 | Yu et al.          | Environ Int                     |
| 88        | 代謝          |                             |           | Early-life exposure to perfluoroalkyl substances in relation to serum adipokines in a longitudinal birth cohort   | 2022 | Shih et al.        | Environ Res                     |
| 89        | 代謝          |                             |           | Association of perfluoroalkyl substances exposure with cardiometabolic traits in an island population of the eastern Adriatic coast of Croatia  | 2019 | Chen et al.        | Sci Total Environ               |
| 90        | 代謝          |                             |           | Perfluoroalkyl substances and metabolic syndrome  | 2019 | Christensen et al. | Int J Hyg Environ Health        |
| 91        | 代謝          |                             |           | Association of serum levels of perfluoroalkyl substances (PFASs) with the metabolic syndrome (MetS) in Chinese male adults: A cross-sectional study   | 2018 | Yang et al.        | Sci Total Environ               |
| 92        | 内分泌・<br>代謝  |                             |           | Positive association between perfluoroalkyl chemicals and hyperuricemia in children   | 2013 | Geiger et al.      | Am J Epidemiol                  |
| 93        | 肝臓          | EPA 2023<br>(PFOS,<br>PFOA) | POD候<br>補 | Liver function biomarkers disorder is associated with<br>exposure to perfluoroalkyl acids in adults: Isomers of<br>C8 Health Project in China   | 2019 | Nian et al.        | Environ Res                     |

| 通し<br>No. | エンドポイント | 評価値算出 機関                    | 区分        | タイトル   | 年    | 著者              | 雑誌                      |
|-----------|---------|-----------------------------|-----------|--|------|-----------------|-------------------------|
| 94        | 肝臓      | EPA 2023<br>(PFOS,<br>PFOA) | POD候<br>補 | Serum perfluorooctanoate (PFOA) and perfluorooctane sulfonate (PFOS) concentrations and liver function biomarkers in a population with elevated PFOA exposure.                                 | 2012 | Gallo et al.    | Environ Health Perspect |
| 95        | 肝臓      | EPA 2023<br>(PFOA)          | POD候<br>補 | Modeled perfluorooctanoic acid (PFOA) exposure and liver function in a mid-Ohio valley community   | 2016 | Darrow et al.   | Environ Health Perspect |
| 96        | 肝臓      |                             |           | Sex differences in the association between perfluoroalkyl acids and liver function in US adolescents: Analyses of NHANES 2013-2016   | 2019 | Attanasio       | Environ Pollut          |
| 97        | 肝臓      |                             |           | Investigation of the Associations Between Low-Dose<br>Serum Perfluorinated Chemicals and Liver Enzymes in<br>US Adults   | 2010 | Lin et al.      | Am J Gastroenterol      |
| 98        | 肝臓      |                             |           | Environmental perfluoroalkyl acid exposures are associated with liver disease characterized by apoptosis and altered serum adipocytokines  | 2019 | Bassler et al.  | Environ Pollut          |
| 99        | 肝臓      |                             |           | Associations of perfluorinated chemical serum concentrations and biomarkers of liver function and uric acid in the US population (NHANES), 2007-2010   | 2015 | Gleason et al.  | Environ Res             |
| 100       | 肝臓      |                             |           | Per- and perfluoroalkyl substances alternatives,<br>mixtures and liver function in adults: A community-<br>based population study in China   | 2022 | Liu et al.      | Environ Int             |
| 101       | 肝臓      |                             |           | Individual and mixture associations of perfluoroalkyl<br>substances on liver function biomarkers in the<br>Canadian Health Measures Survey   | 2022 | Borghese et al. | Environ Health          |
| 102       | 神経      |                             |           | Early-life exposure to persistent organic pollutants<br>(OCPs, PBDEs, PCBs, PFASs) and attention-<br>deficit/hyperactivity disorder: A multi-pollutant<br>analysis of a Norwegian birth cohort | 2019 | Lenters et al.  | Environ Int             |
| 103       | 神経      |                             |           | Childhood exposure to per- and polyfluoroalkyl substances and neurodevelopment in the CHARGE case-control study  | 2022 | Oh et al.       | Environ Res             |
| 104       | 心血管     |                             |           | Association of perfluoroalkyl substances with gestational hypertension and preeclampsia in the MIREC study   | 2020 | Borghese et al. | Environ Int             |
| 105       | 心血管     |                             |           | Serum polyfluoroalkyl chemicals are associated with risk of cardiovascular diseases in national US population  | 2018 | Huang et al.    | Environ Int             |
| 106       | 心血管     |                             |           | Association between perfluoroalkyl acids and the prevalence of hypertension among US adults  | 2020 | Liao et al.     | Ecotoxicol Environ Saf  |
| 107       | 心血管     |                             |           | Perfluorooctanoic acid and cardiovascular disease in US adults   | 2012 | Shankar et al.  | Arch Intern Med         |

| 通し<br>No. | エンドポイント | 評価値算出機関 | 区分 | タイトル  | 年    | 著者              | 雑誌                        |
|-----------|---------|---------|----|---|------|-----------------|---------------------------|
| 108       | 腎臓      |         |    | Perfluoroalkyl substances and kidney function in chronic kidney disease, anemia, and diabetes   | 2018 | Conway et al.   | Diabetes Metab Syndr Obes |
| 109       | 腎臓      |         |    | Renal function and isomers of perfluorooctanoate (PFOA) and perfluorooctanesulfonate (PFOS): Isomers of C8 Health Project in China  | 2019 | Wang et al.     | Chemosphere               |
| 110       | 腎臓      |         |    | Isomers of per- and polyfluoroalkyl substances and uric acid in adults: Isomers of C8 Health Project in China   | 2019 | Zeng et al.     | Environ Int               |
| 111       | 腎臓      |         |    | Perfluoroalkyl substances (PFASs) exposure and kidney damage: Causal interpretation using the US 2003-2018 National Health and Nutrition Examination Survey (NHANES) datasets   | 2021 | Moon et al.     | Environ Pollut            |
| 112       | レビュー    |         |    | Cumulative risk assessment of 17 perfluoroalkylated and polyfluoroalkylated substances (PFASs) in the Swedish population  | 2013 | Borg et al.     | Environ Int               |
| 113       | 方法論     |         |    | Hazard quotient profiles used as a risk assessment tool for PFOS and PFOA serum levels in three distinctive European populations  | 2015 | Ludwicki et al. | Environ Int               |
| 114       | 方法論     |         |    | Derivation of a Human In Vivo Benchmark Dose for Perfluorooctanoic Acid From ToxCast In Vitro Concentration-Response Data Using a Computational Workflow for Probabilistic Quantitative In Vitro to In Vivo Extrapolation         | 2021 | Loizou et al.   | Front Pharmacol           |
| 115       | 分類      |         |    | Commentary: cumulative risk assessment of perfluoroalkyl carboxylic acids and perfluoralkyl sulfonic acids: what is the scientific support for deriving tolerable exposures by assembling 27 PFAS into 1 common assessment group? | 2022 | Colnot et al.   | Arch Toxicol              |

| 通し<br>No. | エンドポイント | 評価値算出 機関           | 区分                       | タイトル   | 年    | 著者                 | 雑誌                              |
|-----------|---------|--------------------|--------------------------|--|------|--------------------|---------------------------------|
| 116       | 発がん性    | EPA 2023<br>(PFOA) | POD                      | Serum concentrations of per- and polyfluoroalkyl substances and risk of renal cell carcinoma   | 2020 | Shearer et al.     | J Natl Cancer Inst              |
| 117       | 発がん性    | EPA 2023<br>(PFOA) |                          | Perfluorooctanoic acid exposure and cancer outcomes in a contaminated community: a geographic analysis.  | 2013 | Vieira et al.      | Environ Health Perspect         |
| 118       | 発がん性    |                    | 専門委<br>員/専<br>門参考<br>人選択 | Perfluorinated alkylated substances serum concentration and breast cancer risk: Evidence from a nested case-control study in the French E3N cohort                                       | 2019 | Mancini et al.     | Int J Cancer                    |
| 119       | 発がん性    |                    | 専門委<br>員/専<br>門参考<br>人選択 | Associations between Polyfluoroalkyl Substances<br>Exposure and Breast Cancer: A Meta-Analysis   | 2022 | Jiang et al.       | Toxics                          |
| 120       | 発がん性    |                    | 専門委<br>員/専<br>門参考<br>人選択 | Plasma perfluoroalkyl substance exposure and incidence risk of breast cancer: A case-cohort study in the Dongfeng-Tongji cohort  | 2022 | Feng et al.        | Environ Pollut                  |
| 121       | 発がん性    |                    |                          | Exposure to perfluoroalkyl substances and risk of hepatocellular carcinoma in a multiethnic cohort   | 2022 | Goodrich et al.    | JHEP Rep                        |
| 122       | 発がん性    |                    |                          | Perfluorooctanoic acid (PFOA) exposures and incident cancers among adults living near a chemical plant   | 2019 | Barry et al.       | Environ Health Perspect         |
| 123       | 発がん性    |                    |                          | Critical Review on PFOA, Kidney Cancer, and<br>Testicular Cancer   | 2020 | Bartell and Vieira | J Air Waste Manag Assoc         |
| 124       | 発がん性    |                    |                          | A critical review of perfluorooctanoate and perfluorooctanesulfonate exposure and cancer risk in humans  | 2014 | Chang et al.       | Crit Rev Toxicol                |
| 125       | 発がん性    |                    |                          | Perfluorooctanoate and perfluorooctanesulfonate plasma levels and risk of cancer in the general Danish population  | 2009 | Eriksen et al.     | J Natl Cancer Inst              |
| 126       | 発がん性    |                    |                          | Glioma is associated with exposure to legacy and alternative per- and polyfluoroalkyl substances   | 2023 | Xie et al.         | J Hazard Mater                  |
| 127       | 発がん性    |                    |                          | Risk assessment for PFOA and kidney cancer based on a pooled analysis of two studies   | 2022 | Steenland et al.   | Environ Int                     |
| 128       | 内分泌     |                    |                          | Serum Biomarkers of Exposure to Perfluoroalkyl<br>Substances in Relation to Serum Testosterone and<br>Measures of Thyroid Function among Adults and<br>Adolescents from NHANES 2011-2012 | 2015 | Lewis et al.       | Int J Environ Res Public Health |
| 129       | 内分泌     |                    |                          | Perfluoroalkyl and polyfluroalkyl substances and maternal thyroid hormones in early pregnancy  | 2020 | Aimuzi et al.      | Environ Pollut                  |

| 通し<br>No. | エンドポイント | 評価値算出 機関 | 区分 | タイトル  | 年    | 著者                       | 雑誌                       |
|-----------|---------|----------|----|---|------|--------------------------|--------------------------|
| 130       | 内分泌     |          |    | Association between organohalogenated pollutants in cord blood and thyroid function in newborns and mothers from Belgian population   | 2018 | Dufour et al.            | Environ Pollut           |
| 131       | 内分泌     |          |    | Pharmacokinetic bias analysis of an association between clinical thyroid disease and two perfluoroalkyl substances  | 2019 | Dzierlenga et al.        | Environ Int              |
| 132       | 内分泌     |          |    | Quantitative bias analysis of the association between subclinical thyroid disease and two perfluoroalkyl substances in a single study   | 2020 | Dzierlenga et al.        | Environ Res              |
| 133       | 内分泌     |          |    | Association between perfluoroalkyl substance exposure and thyroid hormone/thyroid antibody levels in maternal and cord blood: The Hokkaido Study  | 2019 | Itoh et al.              | Environ Int              |
| 134       | 内分泌     |          |    | Association of perfluorinated chemical exposure in utero with maternal and infant thyroid hormone levels in the Sapporo cohort of Hokkaido Study on the Environment and Children's Health                             | 2016 | Kato et al.              | Environ Health Prev Med  |
| 135       | 内分泌     |          |    | Thyroid function and perfluoroalkyl acids in children living near a chemical plant  | 2012 | Lopez-Espinosa<br>et al. | Environ Health Perspect  |
| 136       | 内分泌     |          |    | Association between serum perfluorooctanoic acid (PFOA) and thyroid disease in the U.S. National Health and Nutrition Examination Survey  | 2010 | Melzer et al.            | Environ Health Perspect  |
| 137       | 内分泌     |          |    | Perfluoroalkyl substances and thyroid stimulating hormone levels in a highly exposed population in the Veneto Region  | 2022 | Gallo et al.             | Environ Res              |
| 138       | 内分泌     |          |    | Maternal Perfluoroalkyl Substances, Thyroid<br>Hormones, and DIO Genes: A Spanish Cross-sectional<br>Study  | 2021 | Sarzo et al.             | Environ Sci Technol      |
| 139       | 内分泌     |          |    | Plasma concentrations of per- and polyfluoroalkyl<br>substances at baseline and associations with<br>glycemic indicators and diabetes incidence among<br>high-risk adults in the Diabetes Prevention Program<br>trial | 2017 | Cardenas et al.          | Environ Health Perspect  |
| 140       | 内分泌     |          |    | Perfluoroalkyl substances in older male anglers in<br>Wisconsin   | 2016 | Christensen et al.       | Environ Int              |
| 141       | 内分泌     |          |    | PFOA is associated with diabetes and metabolic alteration in US men: National Health and Nutrition Examination Survey 2003-2012   | 2018 | He et al.                | Sci Total Environ        |
| 142       | 内分泌     |          |    | Nonlinear associations between dietary exposures to perfluorooctanoic acid (PFOA) or perfluorooctane sulfonate (PFOS) and type 2 diabetes risk in women: Findings from the E3N cohort study                           | 2018 | Mancini et al.           | Int J Hyg Environ Health |
| 143       | 内分泌     |          |    | Plasma concentrations of perfluoroalkyl substances and risk of Type 2 diabetes: A prospective investigation among U.S. Women  | 2018 | Sun et al.               | Environ Health Perspect  |

| 通し<br>No. | エンドポイント | 評価値算出 機関 | 区分 | タイトル   | 年    | 著者             | 雑誌                       |
|-----------|---------|----------|----|--|------|----------------|--------------------------|
| 144       | 内分泌     |          |    | Association between per- and polyfluoroalkyl substances and risk of gestational diabetes mellitus  | 2022 | Wang et al.    | Int J Hyg Environ Health |
| 145       | 内分泌     |          |    | Endocrine-disrupting chemicals and the risk of gestational diabetes mellitus: a systematic review and meta-analysis  | 2022 | Yan et al.     | Environ Health           |
| 146       | 内分泌     |          |    | Prenatal exposure to perfluorodecanoic acid is associated with lower circulating concentration of adrenal steroid metabolites during mini puberty in human female infants. The Odense Child Cohort | 2020 | Jensen et al.  | Environ Res              |
| 147       | 内分泌     |          |    | Perfluoroalkyl substances and ovarian hormone concentrations in naturally cycling women  | 2015 | Barrett et al. | Fertil Steril            |
| 148       | その他     |          |    | Associations of serum perfluoroalkyl substance and vitamin D biomarker concentrations in NHANES, 2003-2010   | 2019 | Etzel et al.   | Int J Hyg Environ Health |
| 149       | その他     |          |    | Perfluoroalkyl substances and changes in bone mineral density: A prospective analysis in the POUNDS-LOST study   | 2019 | Hu et al.      | Environ Res              |